



Vascular thrombosis after single dose Ad26.COv2.S vaccine in healthcare workers in South Africa: open label, single arm, phase 3B study (Sisonke study)

Barry Frank Jacobson,¹ Elise Schapkaitz,¹ Azwi Takalani,² Pradeep Rowji,³ Vernon Johan Louw,⁴ Jessica Opie,⁵ Linda-Gail Bekker,⁶ Nigel Garrett,^{7,8} Ameena Goga,^{9,10} Tarylee Reddy,¹¹ Nonhlanhla Yende-Zuma,¹² Ian Sanne,¹³ Ishen Seocharan,¹¹ Jonny Peter,^{14,15} Michelle Robinson,¹⁶ Shirley Collie,¹⁷ Amber Khan ,¹ Simbarashe Takuva,^{18,19} Glenda Gray⁹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjmed-2022-000302>).

For numbered affiliations see end of article.

Correspondence to: Professor Barry Frank Jacobson, Molecular Medicine and Haematology, University of the Witwatersrand Johannesburg Faculty of Health Sciences, Johannesburg, South Africa; clot@nhls.ac.za

Cite this as: *BMJMED* 2023;2:e000302. doi:10.1136/bmjmed-2022-000302

Received: 30 June 2022

ABSTRACT

OBJECTIVE To assess the rates of vascular thrombotic adverse events in the first 35 days after one dose of the Ad26.COv2.S vaccine (Janssen/Johnson & Johnson) in healthcare workers in South Africa and to compare these rates with those observed in the general population.

DESIGN Open label, single arm, phase 3B study.

SETTING Sisonke study, South Africa, 17 February to 15 June 2021.

PARTICIPANTS The Sisonke cohort of 477 234 healthcare workers, aged ≥18 years, who received one dose of the Ad26.COv2.S vaccine.

MAIN OUTCOME MEASURES Observed rates of venous arterial thromboembolism and vaccine induced immune thrombocytopenia and thrombosis

in individuals who were vaccinated, compared with expected rates, based on age and sex specific background rates from the Clinical Practice Research Datalink GOLD database (database of longitudinal routinely collected electronic health records from UK primary care practices using Vision general practice patient management software).

RESULTS Most of the study participants were women (74.9%) and median age was 42 years (interquartile range 33-51). Twenty nine (30.6 per 100 000 person years, 95% confidence interval 20.5 to 44.0) vascular thrombotic events occurred at a median of 14 days (7-29) after vaccination. Of these 29 participants, 93.1% were women, median age 46 (37-55) years, and 51.7% had comorbidities. The observed to expected ratios for cerebral venous sinus thrombosis with thrombocytopenia and pulmonary embolism with thrombocytopenia were 10.6 (95% confidence interval 0.3 to 58.8) and 1.2 (0.1 to 6.5), respectively. Because of the small number of adverse events and wide confidence intervals, no conclusions were drawn between these estimates and the expected incidence rates in the population.

CONCLUSIONS Vaccine induced immune thrombocytopenia and thrombosis after one dose of the Ad26.COv2.S vaccine was found in only a few patients in this South African population of healthcare workers. These findings are reassuring if considered in terms of the beneficial effects of vaccination against covid-19 disease. These data support the continued use of this vaccine, but surveillance is recommended to identify other incidences of venous and arterial thromboembolism and to improve confidence in the data estimates.

TRIAL REGISTRATION ClinicalTrials.gov NCT04838795.

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ In adults, the evidence suggests that the Ad26.COv2.S vaccine (Janssen/Johnson & Johnson) increases the risk of vaccine induced immune thrombocytopenia and thrombosis
- ⇒ Data on the safety of vaccination with Ad26.COv2.S are limited and a direct effect on the development of vaccine induced immune thrombocytopenia and thrombosis has not been established in African populations
- ⇒ Cohort single and multicentre studies, and reviews of vaccine induced immune thrombocytopenia and thrombosis after vaccination with Ad26.COv2.S prompted a cautionary recommendation by the US Centers for Disease Control and Prevention advisory committee that was endorsed by the Food and Drug Administration

WHAT THIS STUDY ADDS

- ⇒ Vaccine induced immune thrombocytopenia and thrombosis was found in only a few participants among South African healthcare workers
- ⇒ The risk of total venous or arterial thromboembolism events was not increased in this large population of healthcare workers who were vaccinated with the Ad26.COv2.S vaccine

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE, OR POLICY

- ⇒ Increased risk of vaccine induced immune thrombocytopenia and thrombosis with the Ad26.COv2.S vaccine in adults emphasises the importance of diagnosis and management in accordance with recommendations
- ⇒ The risk-to-benefit ratio of vaccination with the Ad26.COv2.S vaccine on a population level is encouraging
- ⇒ The Ad26.COv2.S vaccine represents a practical solution for Africa and countries with limited resources, owing to the less stringent storage requirements and single dose scheduling

Introduction

The efficacy of vaccines against SARS-CoV-2 has been reported to be 53-97% in patients with symptoms of infection.¹⁻⁶ Of these, the adenovirus vector vaccines are the preferred vaccines in countries with limited resources because of their accessibility and less stringent storage requirements. Two of these include the chimpanzee

adenoviral vector, ChAdOx1 nCoV-19 vaccine (Oxford-AstraZeneca) and the adenoviral vector type 26, Ad26.COV2.S vaccine (Janssen/Johnson & Johnson). The risk of adverse events after these vaccines has been assessed in clinical trials and by public health surveillance systems. Reports of serious adverse events have been rare. Nonetheless, some instances of venous thromboembolism and arterial thrombosis after adenoviral vector vaccines have been reported, raising concerns about the potential risks of thromboembolic complications.⁷⁻¹⁰

Moderate-to-severe thrombocytopenia and thrombosis at unusual sites have been reported in rare instances after vaccination with the ChAdOx1 nCoV-19 and Ad26.COV2.S vaccines.¹¹⁻¹² This finding prompted a recommendation by several regulatory authorities for a temporary suspension of vaccinations in April 2021.¹³ The term vaccine induced immune thrombocytopenia and thrombosis was created for this syndrome, with a proposed mechanism similar to that of heparin induced thrombocytopenia with thrombosis.¹⁴⁻¹⁵ Also, the term thrombosis with thrombocytopenia syndrome has been defined by the Brighton Collaboration.¹⁶ Thrombosis with thrombocytopenia syndrome can be attributed to several causes and the term is recommended for use in epidemiological studies after vaccination.

Definite or probable vaccine induced immune thrombocytopenia and thrombosis was reported in 220 patients presenting to hospitals in the UK after 24 million first doses of the ChAdOx1 nCoV-19 vaccine.¹⁷ Median age of these patients was 48 years (range 18-79) and 55% were women. Of these, 209 (95%) presented with thrombocytopenia (median $47 \times 10^9/L$, range 28-76) and 102 (46%) with cerebral venous sinus thrombosis. No other risk factors for thrombosis were identified and the overall mortality rate was 22%. Up to 8 July 2021, another 38 patients with definite vaccine induced immune thrombocytopenia and thrombosis after 12.5 million first doses of Ad26.COV2.S vaccine were reported to the US Centers for Disease Control and Prevention advisory committee on immunisation practices.¹⁸ These patients had a similar presentation to those described after vaccination with the ChAdOx1 nCoV-19 vaccine, suggesting an effect of adenoviral vector-type vaccines against SARS-CoV-2. Thrombosis at unusual sites and thrombocytopenia have also been reported for mRNA based vaccines, although to a lesser extent.¹⁰⁻¹⁹ Given the severity of vaccine induced immune thrombocytopenia and thrombosis and the ongoing covid-19 pandemic, close monitoring and reporting of serious adverse events is essential.

Between 17 February and 17 May 2021, the Ad26.COV2.S vaccine was given to 477 234 individuals in the Sisonke study, a multicentre, open label, single arm, phase 3B implementation study in healthcare workers in South Africa. Preliminary data on vascular thrombotic events after vaccination of the first 288 368 healthcare workers have been reported.²⁰ The rate of venous and arterial thromboembolic events were 1.7 per 100 000 participants. These events occurred in individuals with known risk factors for thromboembolism. Here, we describe the occurrence of vascular thrombotic events after vaccination of 477 234 healthcare workers in the Sisonke study, including two patients with vaccine induced immune thrombocytopenia and thrombosis.

Methods

Study participants and procedures

The Sisonke study has been previously described.²¹ After the temporary suspension of vaccinations in South Africa on 13 April 2021, vaccinations resumed on 28 April 2021, with further safeguards. These safeguards included: changes to the eligibility criteria in the Sisonke protocol; more information for participants about the signs and symptoms of thrombosis; amendments to patient information sheets and informed consent forms; and modified standard operating procedures with training to support appropriate screening as well as management of participants at high risk of thrombosis. Also, a short message surveillance service was implemented with specific increased awareness of the risk of thrombosis.

Vaccinations were conducted in collaboration with the South African Department of Health in public or private vaccination centres nationwide. Delivery of the vaccine was managed by appropriately trained personnel linked to the ENSEMBLE (A Study of Ad26.COV2.S for the Prevention of SARS-CoV-2-Mediated COVID-19 in Adult Participants) trial research sites (ClinicalTrials.gov NCT04505722). All healthcare workers, registered with the Electronic Vaccination Data System, the South African national vaccination registry, and who provided consent for the study electronically were eligible for enrolment. The study excluded women who were breastfeeding (after 28 April 2021) and pregnant women because of lack of sufficient safety data.

Participants received one intramuscular injection of Ad26.COV2.S at a dose of 5×10^{10} virus particles, and were observed for adverse events for 15 minutes after vaccination, or for 30 minutes if they had a previous history of allergic reactions to vaccinations. The Sisonke study was a single dose study. A second dose study, Sisonke, 2 was performed after findings from international




Figure 1 | Flowchart of study participants

studies showed the protective effect of a second dose.²²

Safety monitoring

Adverse events in the Sisonke study were reported from a hybrid surveillance system that combined passive and active reporting of adverse event outcomes. Full details of the Sisonke pharmacovigilance and safety reporting processes are detailed elsewhere.²¹ Adverse event reports were reviewed and processed daily by the safety team, and validated and updated by telephonic interviews with participants, healthcare providers, and from accessible medical records. A safety review team of public health experts and specialists in neurology and haematology met weekly to review the cumulative safety data and case summaries. An independent safety monitoring committee provided safety oversight. In this analysis, we evaluated all data up to 15 June 2021 (35 days after the last vaccination). Surveillance for adverse event is ongoing and will continue for up to two years after vaccination.

Case definitions

Safety events of concern included adverse events of special interest: vascular thrombosis after vaccination, based on the Brighton Collaboration list, and serious adverse events after vaccination that met the criteria of the International Conference on Harmonisation.²³ Vascular thrombosis included one or more episodes of venous, arterial, or small vessel thrombosis. Venous thromboembolism events were radiologically confirmed by cerebral magnetic resonance venography and compression ultrasound of the limbs, with or without computed tomography angiography of the chest and abdomen.

Arterial thrombosis events included acute coronary syndrome diagnosed by electrocardiogram and cardiac biomarkers, and ischaemic stroke diagnosed by neurological examination in combination with computed tomography or magnetic resonance imaging. Small vessel thrombosis was diagnosed by histopathology in any tissue or organ.

Definite vaccine induced immune thrombocytopenia and thrombosis was defined as: onset of symptoms 5-30 days after vaccination, presence of thrombosis, platelet count $<150 \times 10^9/L$ or decrease of 50% from baseline, and presence of antibodies to platelet factor 4.^{24 25} The diagnosis was further supported by plasma concentrations of d-dimers >0.25 mg/L and fibrinogen <2 g/L. If two of the four criteria were met, participants were defined as having possible vaccine induced immune thrombocytopenia and thrombosis (online supplemental tables 1S and 2S).¹⁷

Blood sampling and laboratory methods

Testing for the SARS-CoV-2 virus was performed in all participants with a real time polymerase chain reaction test of the upper respiratory tract (nasopharyngeal and oropharyngeal swab). As far as possible, serological testing was performed. Antibodies to platelet factor 4 were detected by commercial platelet factor 4/heparin enzyme linked immunoassays (Zymutest HIA IgG; Hyphen BioMed, Neuville-sur-Oise, France).²⁵ The positive threshold was based on the manufacturer's optical density threshold (reference range <0.30 negative, 0.3-0.5 weak positive, and >0.5 strong positive). Positive results were confirmed by a functional assay^{26 27} (platelet aggregation profiler, PAP-8E, reference range $<25\%$; Bio/Data Corporation, Horsham, PA) at a centralised referral laboratory.

Data analysis

For all data transfer during the study, industry standard multilayer encryption platforms were used. The Electronic Vaccination Data System used a beneficiary identifier, a unique 36 character machine generated identifier within the system. This identifier was used for data linkages, removing any direct identifiers.

Statistical analyses were conducted with Stata, version 14. For descriptive statistics, frequencies and percentages were used for categorical variables and median (interquartile range) for continuous variables. Occurrence of venous and arterial thromboembolic events were described by baseline characteristics of participants. The observed number of reported incidences was compared with the expected number based on background incidence rates, and the observed to expected ratio (with 95% confidence interval) was calculated. The expected number of events was estimated based on background incidence rates of vascular

thrombotic events in the Clinical Practice Research Datalink GOLD database (database of longitudinal routinely collected electronic health records from UK primary care practices using Vision general practice patient management software).^{28 29} This database contains data from 3 913 071 participants with a median age of 41 years (interquartile range 22-59), contributed by general practitioners from the UK between 1 January 2017 and 31 December 2019. These background incidence rates were applied because no data were available locally. The Clinical Practice Research Datalink GOLD database was selected because it includes thromboses of special interest for covid-19 vaccinations, with rates grouped by sex and age.³⁰ Person time was accrued from the date of vaccination until death or closure of the dataset on 15 June 2021 (35 days after the last vaccination). The incidence rates per 100 000 person years were calculated with a Poisson model, with person years as an offset.

Patient and public involvement

No patients were involved in the design, execution, or interpretation of this study. Owing to the urgency and sensitivity of the study question, as well as data privacy constraints, involving members of the public in the study was not possible. The results of the study have been disseminated to research participants and public

communities electronically as well as via the press, parliamentary committees, and annual reports.

Results

Participant characteristics

The Sisonke cohort included 119 753 (25.1%) male participants and 357 481 (74.9%) female participants, with a median age of 42 (interquartile range 33-51) years. Data for sex were taken from information in the Sisonke study rather than from patient reported gender. The higher proportion of women was because the nursing profession in South Africa has disproportionately more women than men. Vascular thrombotic events were reported for 37 participants. Eight participants were excluded (figure 1) and 29 (30.6 per 100 000 person years, 95% confidence interval 20.5 to 44.0) vascular thrombotic events were included. Table 1 shows the baseline characteristics of the 29 participants. Most participants were women (n=27, 93.1%) and median age was 46 years (interquartile range 37-55). Median time from vaccination to onset of thrombosis was 14 days (7-29). Baseline characteristics of participants who presented with venous thromboembolism, arterial thrombosis, or vaccine induced immune thrombocytopenia and thrombosis were not substantially different.

Table 1 | Baseline characteristics of participants with venous thromboembolism, arterial thrombosis, and vaccine induced immune thrombocytopenia and thrombosis in the Sisonke study

Characteristics	Sisonke study	All thrombotic events	Venous thromboembolism	Arterial thrombosis	Vaccine induced immune thrombocytopenia and thrombosis
Total No of participants	477 234	29 (100)	18 (62.1)	9 (31.0)	2 (6.9)
Sex:					
Men	119 753 (25.1)	2 (6.9)	1 (5.6)	1 (11.1)	0 (0)
Women	357 481 (74.9)	27 (93.1)	17 (94.4)	8 (88.9)	2 (100.0)
Age group (years):					
18-39	209 411 (43.9)	10 (34.5)	7 (38.9)	2 (22.2)	1 (50.0)
40-49	136 967 (28.7)	10 (34.5)	6 (33.3)	3 (22.2)	1 (50.0)
50-59	96 235 (20.2)	3 (10.3)	0 (0.0)	3 (33.3)	0 (0.0)
≥60	23 389 (7.2)	6 (20.7)	5 (27.8%)	1 (11.1%)	0 (0.0)
Median (IQR) age (years)	42 (33-51)	46 (37-55)	44 (34-60)	46 (43-55)	39 (28-49)
Median (IQR) time to onset of thrombosis (days)	–	14 (7-29)	16 (7-23)	10 (5-38)	19 (7-30)
Comorbidities:					
Cancer	1 364 (0.3)	0 (0)	0 (0)	0 (0)	0 (0)
Chronic lung disease	1 733 (0.4)	0 (0)	1 (5.6)	0 (0)	0 (0)
Diabetes	28 063 (5.9)	3 (10.3)	1 (5.6)	0 (0)	0 (0)
Heart disease	3 430 (0.7)	2 (6.9)	0 (0)	1 (8.3)	0 (0)
HIV	39 386 (8.3)	2 (6.9)	1 (5.6)	1 (8.3)	0 (0)
Hypertension	74 381 (15.6)	8 (27.6)	3 (16.7)	3 (33.3)	0 (0)
Tuberculosis	478 (0.1)	0 (0)	0 (0)	0 (0)	0 (0)

Data are number (%) unless stated otherwise.

IQR=interquartile range.




Figure 2 | Categories of vascular thrombosis. Other arterial thrombosis included cephalic vein thrombosis (n=1) and retinal vein thrombosis (n=1)

Description of vascular thrombotic events

Eighteen (62.1%) venous thromboembolism events were observed. Of these, pulmonary embolism and deep vein thrombosis were the most common venous events, affecting 15 (83.3%) of the 18 participants (figure 2). Other venous sites were uncommon presentations: cephalic vein thrombosis (n=1, 5.6%) and retinal vein thrombosis (n=1, 5.6%). One (5.6%) cerebral venous sinus thrombosis event was reported in a woman (aged 30-39 years). The woman presented with a persistent headache followed by a seizure 29 days after vaccination, associated with a normal platelet count and negative antiplatelet factor 4 antibodies. When the analysis was grouped by age and sex, the observed to expected ratio for cerebral venous sinus thrombosis was 2.8 (95% confidence interval 0.1 to 15.4) for women aged 30-39 years.

Of the nine (31.0%) arterial thrombotic events, five (55.6%) were ischaemic strokes, two (22.2%) were acute coronary syndrome, and one (11.1%) was an ischaemic limb associated with positive antiphospholipid antibodies. Also, one patient (11.1%) had leucocytoclastic vasculitis, diagnosed by histopathological examination of a biopsy sample. We found no evidence of a clear association between previous vaccination with the Ad26.COV2.S vaccine and arterial thrombosis after vaccination. When the analysis was grouped by age and sex, the observed to expected ratio for ischaemic strokes was 2.5 (95% confidence interval 0.3 to 9.0) for women aged 30-39 years and 1.2 (0.1 to 4.2) for women aged 40-49 years.

Of the 29 participants who presented with vascular thrombotic events, two (6.9%) fulfilled the diagnostic criteria for definite or possible vaccine induced immune thrombocytopenia and thrombosis; the first

had a pulmonary embolus with thrombocytopenia and the second a cerebral sinus venous thrombosis with thrombocytopenia. These results corresponded to an observed to expected ratio of 1.2 (95% confidence interval 0.1 to 6.5) and 10.6 (0.3 to 58.8), respectively. The first participant was a woman (aged 40-49 years) who presented with a saddle pulmonary embolism, platelet count of $34 \times 10^9/L$, positive antiplatelet factor 4 antibodies of 0.86, confirmed by a functional heparin induced thrombocytopenia with thrombosis assay (>80%), and plasma concentrations of d-dimers >10.0 mg/L, nine days after vaccination. The second participant was a woman (aged 20-29 years) who was admitted to hospital unconscious after a severe headache, restlessness, and confusion, one month after vaccination. A computed tomography venogram was consistent with cerebral venous sinus thrombosis. The antiplatelet factor 4 antibody assay was negative, and she had marginal thrombocytopenia ($146 \times 10^9/L$). Both participants recovered.

At the time of the data analysis, two deaths (6.9%) related to vascular thrombosis were reported. The first participant was a woman (aged 60-69 years) with a previous history of venous thromboembolism, who collapsed 23 days after vaccination. A pulmonary embolism was diagnosed on post-mortem examination. The second participant was a woman (aged 30-39 years) who presented with a pulmonary embolism, 41 days after vaccination. She died after admission to hospital, 42 days after vaccination.

Risk factors for vascular thrombotic events

Of the 29 vascular events, 15 (51.7%) occurred in participants with one or more comorbidities (table 2), and four (13.8%) participants had a history of thrombosis.

Discussion

Principal findings

Safety concerns about vascular thrombotic events have been raised after vaccination with adenoviral vector vaccines against SARS-CoV-2 infection. Venous thromboembolism, arterial thrombosis, and vaccine induced immune thrombocytopenia and thrombosis events were observed after the first dose of the Ad26.COV2.S vaccine. This finding prompted a cautionary recommendation by the US Centers for Disease Control and Prevention advisory committee which was endorsed by the Food and Drug Administration. This South African population based study of almost half a million healthcare workers, aged ≥ 18 years, who received the Ad26.COV2.S vaccine, provides further real world evidence that the risk of vascular thrombotic events seems to be low, supporting its continued use in this setting.

Table 2 | Observed versus expected vascular thrombotic events after vaccination with the Ad26.COV2.S vaccine in the Sisonke study, by sex and age

	Observed count	Person years	Observed incidence rate per 100 000 person years (95% CI)	Expected incidence rate per 100 000 person years (95% CI)	Expected count	Observed to expected ratio (95% CI)
Venous thromboembolism						
Total No of pulmonary embolism events	11	94 743.07	11.61 (6.43 to 20.97)	74.0 (72.3 to 75.7)	70.11	0.16 (0.08 to 0.28)
Men:						
Aged 60-69 years	1	1677.95	59.60 (8.39 to 423.08)	151.3 (141.1 to 162.1)	2.54	0.39 (0.01 to 2.19)
Women:						
Aged 30-39 years	3	20 135.48	14.90 (4.81 to 46.20)	40.1 (35.4 to 45.2)	8.07	0.37 (0.08 to 1.09)
Aged 40-49 years	4	21 204.73	18.86 (7.08 to 50.26)	50.1 (44.9 to 55.8)	10.62	0.38 (0.10 to 0.96)
Aged 60-69 years	3	4137.99	72.50 (23.38 to 224.79)	134.9 (125.4 to 145.0)	5.58	0.54 (0.11 to 1.57)
Total No of deep vein thrombosis events	4	94 743.07	4.22 (1.58 to 11.25)	93.9 (92.0 to 95.9)	88.96	0.04 (0.01 to 0.12)
Women:						
Aged 20-29 years	1	9989.95	10.01 (1.41 to 71.06)	58.5 (52.4 to 65.2)	5.84	0.17 (0 to 0.95)
Aged 30-39 years	2	20 135.48	9.93 (2.48 to 39.72)	71.8 (65.5 to 78.6)	14.46	0.14 (0.02 to 0.5)
Aged 40-49 years	1	21 204.73	4.72 (0.66 to 33.48)	80.4 (73.7 to 87.5)	17.05	0.06 (0 to 0.33)
Total No of cerebral venous sinus thrombosis events	1	94 743.07	1.06 (0.15 to 7.49)	1.2 (1.0 to 1.5)	1.14	0.88 (0.02 to 4.9)
Women:						
Aged 30-39 years	1	20 135.48	4.97 (0.70 to 35.26)	1.8 (0.9 to 3.2)	0.36	2.76 (0.07 to 15.37)
Arterial thrombosis						
Total No of ischaemic stroke events	5	94 743.07	5.28 (2.20 to 12.68)	29.6 (28.5 to 30.7)	28.04	0.18 (0.06 to 0.42)
Women:						
Aged 30-39 years	2	20 135.48	9.93 (2.48 to 39.72)	4.0 (2.6 to 5.8)	0.81	2.48 (0.30 to 8.97)
Aged 40-49 years	2	21 204.73	9.43 (2.36 to 37.71)	8.2 (6.2 to 10.7)	1.74	1.15 (0.14 to 4.16)
Aged 50-59 years	1	15 054.54	6.64 (0.94 to 47.16)	17.3 (14.4 to 20.7)	2.6	0.38 (0.01 to 2.14)
Total No of acute coronary syndrome events	2	94 743.07	2.11 (0.53 to 8.44)	138.8 (136.5 to 141.2)	131.5	0.02 (0 to 0.05)
Men:						
Aged 60-69 years	1	1677.95	59.60 (8.39 to 423.08)	437.2 (419.7 to 455.4)	7.34	0.14 (0 to 0.76)
Women:						
Aged 40-49 years	1	21 204.73	4.72 (0.66 to 33.48)	33.9 (29.7 to 38.7)	7.19	0.14 (0 to 0.78)
Vaccine induced immune thrombocytopenia and thrombosis						
Total No of pulmonary embolism and thrombocytopenia events	1	94 743.07	1.06 (0.15 to 7.49)	0.9 (0.7 to 1.1)	0.85	1.17 (0.03 to 6.53)
Women:						
Aged 40-49 years	1	21 204.73	4.72 (0.66 to 33.48)	0.3 (0.1 to 0.8)	0.06	15.72 (0.4 to 87.58)
Total No of cerebral sinus venous thrombosis and thrombocytopenia events	1	94 743.07	1.06 (0.15 to 7.49)	0.1 (0.1 to 0.2)	0.09	10.55 (0.27 to 58.81)
Women:						
Aged 20-29 years	1	9989.98	10.01 (1.41 to 71.06)	–	–	–

CI=confidence interval.

Comparison with other studies

Venous thromboembolism is among the most common causes of morbidity and mortality in Africa.³¹ When considering the effect of vaccination on public health, the background incidence rate of thrombosis in the general population, as well as according to age and sex, should be taken into

account. Consistent with other population based studies, the most common thrombotic event in this study population was venous thromboembolism.⁷⁻⁹ The rates for total venous thromboembolism events were below the expected background incidence rates of the Clinical Practice Research Datalink GOLD database. This finding is in contrast with the

substantially increased rates of venous thromboembolism secondary to covid-19 disease.^{32–34} In particular, earlier strains of the SARS-CoV-2 virus were associated with an increased rate of pulmonary embolism (>30%) in patients admitted to hospital.^{35–37} Nonetheless, we found an increase in the incidence of definite or possible vaccine induced immune thrombocytopenia and thrombosis. Vaccine induced immune thrombocytopenia and thrombosis has consistently been reported as a rare event in the general population.^{11 12} These two (6.9%) patients (pulmonary embolus with thrombocytopenia and cerebral sinus venous thrombosis with thrombocytopenia) were diagnosed and managed with the support of the protocol safety review team, and both participants recovered. Timely identification of these events will ensure the successful management of vaccine induced immune thrombocytopenia and thrombosis according to published recommendations.²⁵

Furthermore, we found no increase in the rates of arterial thromboembolism. Our subanalysis according to age and sex, in women aged 30–39 years, showed three ischaemic strokes in the vaccine cohort compared with one expected event. This finding should be interpreted in terms of recently published data which reported a 2–3-fold higher incidence of stroke in Africa than in Europe.³⁸ Also, more than half of the vascular thrombotic events in our study occurred in participants with one or more cardiovascular risk factors. Moreover, according to a recent meta-analysis, patients with severe covid-19 disease have a fivefold increased risk of stroke.³⁹ Specific immune mediated and prothrombotic mechanisms contribute to the increased risk of vascular thrombotic events, providing another clear reason for vaccination.

Strengths and limitations of this study

The findings of the Sisonke study have important regulatory implications. The Ad26.COVS vaccine showed >60% protection against severe covid-19 disease and >80% protection against death related to SARS-CoV-2 infection.⁴⁰ Therefore, the risk-to-benefit ratio of vaccination at the population level is encouraging. From a public health perspective, the Ad26.COVS vaccine represents a practical solution for Africa, with limited resources,²² owing to the less stringent storage requirements and single dose scheduling associated with reasonable efficacy.⁴⁰

Our study had several limitations. Firstly, the Sisonke study is mainly a passive surveillance system, relying on self-reporting, and therefore adverse events after vaccination with the Ad26.COVS vaccine might have been under-reported. Because the study cohort were healthcare workers, however, adverse events would likely have been reported, with the the potential for overreporting of adverse events. Also, the temporary suspension in adenovirus based vaccinations in April 2021 could have increased participants' levels of

clinical awareness and self-reporting of adverse events. In our study, comparison of adverse events with mRNA based vaccines was not possible because Ad26.COVS was the only available vaccine at the time of this study. In the study of Cari et al,⁶ to reduce under-reporting and overreporting biases, adverse events with a mRNA based covid-19 vaccine were compared with adenovirus based vaccines. All adverse events assessed in this study were verified by our study investigators.

Secondly, the observed to expected analysis should be considered in terms of the non-South African reference data. The background incidence rates of the GOLD database were selected because they included thromboses of special interest for covid-19 vaccinations, with rates grouped by sex and age. Subsequently, Discovery Health conducted a study of South African medical insurance claims for adverse events related to the Ad26.COVS vaccine between February 2021 and September 2021. This study compared adverse events (venous thromboembolic events, stroke, and myocardial infarction) in 103 315 recipients of a vaccine and matched individuals who were not vaccinated (online supplemental table 4) but no significant differences were found.

Thirdly, owing to the small number of adverse events, comparisons with the incidence in the expected population were associated with wide 95% confidence intervals and therefore no conclusions could be drawn from the data. Ongoing surveillance is needed to identify further venous and arterial thromboembolism events and to improve the accuracy of the data estimates. Fourthly, the Sisonke study did not report participants' race and therefore adverse events could not be grouped by race. Lastly, risk factors for venous thromboembolism, such as history of thrombosis, inherited thrombophilia, and use of oral contraceptives, were not available for all participants.

Conclusions

In our pharmacovigilance study of single dose Ad26.COVS vaccine, we found two patients with vaccine induced immune thrombocytopenia and thrombosis among almost half a million recipients. The risks of total venous or arterial thromboembolism events were also low. The incidence of cerebral and splanchnic venous thrombosis with or without thrombocytopenia seemed to be lower after vaccination with the Ad26.COVS vaccine than the ChAdOx1 nCoV-19 vaccine, as reported by the EudraVigilance European database.¹⁹ These risks, when interpreted with data on the effectiveness of the Ad26-CoV2.S vaccine against severe disease and death, emphasise the benefit versus risk of the Ad26.CoV2.S vaccine at the population level.

AUTHOR AFFILIATIONS

¹Molecular Medicine and Haematology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

- ²Hutchinson Centre Research Institute of South Africa (HCRISA), Chris Hani Baragwanath Hospital, Johannesburg, South Africa
- ³The Southern African Society of Thrombosis and Haemostasis, Neurology Association of South Africa, Johannesburg, South Africa
- ⁴Division of Clinical Haematology, Department of Medicine, Faculty of Health Sciences, University of Cape Town and Groote Schuur Hospital, Cape Town, South Africa
- ⁵Division of Haematology, Department of Pathology, University of Cape Town and National Health Laboratory Service, Cape Town, South Africa
- ⁶Desmond Tutu HIV Centre, University of Cape Town, Cape Town, South Africa
- ⁷Centre for the AIDS Programme of Research in South Africa, Durban, South Africa
- ⁸Discipline of Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, Durban, South Africa
- ⁹HIV and Other Infectious Diseases Research Unit, South African Medical Research Council, Cape Town, South Africa
- ¹⁰Paediatrics and Child Health, University of Pretoria, Pretoria, South Africa
- ¹¹Biostatistics Research Unit, South African Medical Research Council, Durban, South Africa
- ¹²Nelson R Mandela School of Medicine, Centre for the AIDS Programme of Research in South Africa, University of KwaZulu Natal, Durban, South Africa
- ¹³Clinical HIV Research Unit, University of the Witwatersrand Faculty of Sciences, Johannesburg, South Africa
- ¹⁴Division of Allergy and Clinical Immunology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa
- ¹⁵Allergy and Immunology Unit, University of Cape Town Lung Institute, Cape Town, South Africa
- ¹⁶Right To Care, Johannesburg, South Africa
- ¹⁷Discovery Health, Johannesburg, South Africa
- ¹⁸School of Health Systems and Public Health, University of Pretoria, Faculty of Health Sciences, Pretoria, South Africa
- ¹⁹Perinatal HIV Research Unit, University of the Witwatersrand, Faculty of Health Sciences, Johannesburg, South Africa

Correction notice This article has been corrected since it first published. Author affiliations have been corrected.

Acknowledgements We thank Tommy Chen, Dave Jacobs, Donald Ntjana, Jared Champion, Karl Bernhardt, and Smybbin Mathews for their insightful contribution to the manuscript.

Contributors ST, AT, ISe, NYZ, ISa, LGB, GG, AG, and NG conceived the study. ST, AT, JP, JO, VL, PFR, BFJ, ISa, LGB, MR, GG, AG, and NG were members of the safety team. The data were curated by ST, AT, ISe, NYZ, TR, AG, and NG. The data were analysed by NYZ, TR, and SC. ES, ST, and BFJ wrote the original manuscript. All authors critically reviewed the manuscript and all revisions. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. BFJ is the guarantor. Transparency: The lead author (the guarantor) affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

Funding The study was funded by the National Treasury of South Africa, National Department of Health, Solidarity Response Fund NPC, Michael and Susan Dell Foundation, Elma Vaccines and Immunisation Foundation, and Bill and Melinda Gates Foundation. The funders had no role in considering the study design or in the collection, analysis, interpretation of data, writing of the report, or decision to submit the article for publication.

Competing interests All authors have completed the ICMJE uniform disclosure form at www.icmje.org/disclosure-of-interest/ and declare: support from National Treasury of South Africa, National Department of Health, Solidarity Response Fund NPC, Michael and Susan Dell Foundation, Elma Vaccines and Immunisation Foundation, and Bill and Melinda Gates Foundation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Ethics approval This study involves human participants and was approved by the South African Health Products Regulatory Authority (reference 20200465). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Amber Khan <http://orcid.org/0000-0002-9950-0464>

REFERENCES

- Sadoff J, Gray G, Vandebosch A, *et al*. Final analysis of efficacy and safety of single-dose Ad26.COV2.S. *N Engl J Med* 2022;386:847–60. [10.1056/NEJMoa2117608](https://doi.org/10.1056/NEJMoa2117608)
- Haas EJ, Angulo FJ, McLaughlin JM, *et al*. Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data. *Lancet* 2021;397:1819–29. [10.1016/S0140-6736\(21\)00947-8](https://doi.org/10.1016/S0140-6736(21)00947-8)
- Voysey M, Clemens SAC, Madhi SA, *et al*. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet* 2021;397:99–111. [10.1016/S0140-6736\(20\)32661-1](https://doi.org/10.1016/S0140-6736(20)32661-1)
- El Sahly HM, Baden LR, Essink B, *et al*. Efficacy of the mRNA-1273 SARS-CoV-2 vaccine at completion of blinded phase. *N Engl J Med* 2021;385:1774–85. [10.1056/NEJMoa2113017](https://doi.org/10.1056/NEJMoa2113017)
- Polack FP, Thomas SJ, Kitchin N, *et al*. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med* 2020;383:2603–15. [10.1056/NEJMoa2034577](https://doi.org/10.1056/NEJMoa2034577)
- Cari L, Alhosseini MN, Fiore P, *et al*. Cardiovascular, neurological, and pulmonary events following vaccination with the BNT162b2, ChAdOx1 nCoV-19, and Ad26.COV2.S vaccines: an analysis of european data. *J Autoimmun* 2021;125:102742. [10.1016/j.jaut.2021.102742](https://doi.org/10.1016/j.jaut.2021.102742)
- European Medicines Agency. COVID-19 vaccine safety update. *vaxzevria* 2021. Amsterdam, The Netherlands: EMA, 2021.
- European Medicines Agency. News: astrazeneca's COVID-19 vaccine: EMA finds possible link to very rare cases of unusual blood clots with low blood platelets. Amsterdam, The Netherlands: EMA, 2021.
- Smadja DM, Yue Q-Y, Chocron R, *et al*. Vaccination against COVID-19: insight from arterial and venous thrombosis occurrence using data from vigibase. *Eur Respir J* 2021;58:2100956. [10.1183/13993003.00956-2021](https://doi.org/10.1183/13993003.00956-2021)
- Cari L, Fiore P, Naghavi Alhosseini M, *et al*. Blood clots and bleeding events following BNT162b2 and ChAdOx1 nCoV-19 vaccine: an analysis of european data. *J Autoimmun* 2021;122:102685. [10.1016/j.jaut.2021.102685](https://doi.org/10.1016/j.jaut.2021.102685)
- Schultz NH, Sørvoll IH, Michelsen AE, *et al*. Thrombosis and thrombocytopenia after ChAdOx1 nCoV-19 vaccination. *N Engl J Med* 2021;384:2124–30. [10.1056/NEJMoa2104882](https://doi.org/10.1056/NEJMoa2104882)
- See I, Su JR, Lale A, *et al*. US case reports of cerebral venous sinus thrombosis with thrombocytopenia after Ad26.COV2.S vaccination, March 2 to April 21, 2021. *JAMA* 2021;325:2448–56. [10.1001/jama.2021.7517](https://doi.org/10.1001/jama.2021.7517)
- Wise J. Covid-19: european countries suspend use of oxford-astrazeneca vaccine after reports of blood clots. *BMJ* 2021;372:699. [10.1136/bmj.n699](https://doi.org/10.1136/bmj.n699)
- Greinacher A, Thiele T, Warkentin TE, *et al*. Thrombotic thrombocytopenia after chadox1 nCoV-19 vaccination. *N Engl J Med* 2021;384:2092–101. [10.1056/NEJMoa2104840](https://doi.org/10.1056/NEJMoa2104840)

- 15 Scully M, Singh D, Lown R, *et al.* Pathologic antibodies to platelet factor 4 after ChAdOx1 nCoV-19 vaccination. *N Engl J Med* 2021;384:2202–11. 10.1056/NEJMoa2105385
- 16 Brighton Collaboration. Interim case definition of thrombosis with thrombocytopenia syndrome (TTS). 2021. Available: <https://brightoncollaboration.us/thrombosis-with-thrombocytopenia-syndrome-interim-case-definition/Date> [Accessed 23 Sep 2022].
- 17 Pavord S, Scully M, Hunt BJ, *et al.* Clinical features of vaccine-induced immune thrombocytopenia and thrombosis. *N Engl J Med* 2021;385:1680–9. 10.1056/NEJMoa2109908
- 18 MacNeil JR, Su JR, Broder KR, *et al.* Updated recommendations from the Advisory Committee on immunization practices for use of the janssen (Johnson & Johnson) COVID-19 vaccine after reports of thrombosis with thrombocytopenia syndrome among vaccine recipients—United States, April 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:651–6. 10.15585/mmwr.mm7017e4
- 19 European Medicines Agency. EudraVigilance: European database of suspected adverse drug reaction reports. Amsterdam, The Netherlands: EMA. 2021. Available: www.adrreports.eu/en/background.html [Accessed Apr 2021].
- 20 Takuva S, Takalani A, Garrett N, *et al.* Thromboembolic events in the south african ad26.COV2.S vaccine study. *N Engl J Med* 2021;385:570–1. 10.1056/NEJMc2107920
- 21 Takuva S, Takalani A, Seocharan I, *et al.* Safety evaluation of the single-dose ad26.COV2.S vaccine among healthcare workers in the sisonke study in South Africa: a phase 3B implementation trial. *PLoS Med* 2022;19:e1004024. 10.1371/journal.pmed.1004024
- 22 Bonhoeffer J, Kohl K, Chen R, *et al.* The Brighton collaboration: addressing the need for standardized case definitions of adverse events following immunization (AEFI). *Vaccine* 2002;21:298–302. 10.1016/S0264-410X(02)00449-8
- 23 Solomon CG, Greinacher A. Heparin-Induced thrombocytopenia. *N Engl J Med* 2015;373:252–61. 10.1056/NEJMcpr1411910
- 24 Jacobson BF, Schapkaitz E, Mer M, *et al.* Recommendations for the diagnosis and management of vaccine-induced immune thrombotic thrombocytopenia. *S Afr Med J* 2021;111:535–7.
- 25 Sheridan D, Carter C, Kelton JG. A diagnostic test for heparin-induced thrombocytopenia. *Blood* 1986;67:27–30. 10.1182/blood.V67.1.27.27
- 26 Nazy I, Sachs UJ, Arnold DM, *et al.* Recommendations for the clinical and laboratory diagnosis of VITT against COVID-19: communication from the ISTH SSC Subcommittee on platelet immunology. *J Thromb Haemost* 2021;19:1585–8. 10.1111/jth.15341
- 27 Herrett E, Gallagher AM, Bhaskaran K, *et al.* Data resource profile: clinical practice research Datalink (CPRD). *Int J Epidemiol* 2015;44:827–36. 10.1093/ije/dyv098
- 28 Wolf A, Dedman D, Campbell J, *et al.* Data resource profile: clinical practice research Datalink (CPRD) aum. *Int J Epidemiol* 2019;48:1740–1740g. 10.1093/ije/dyzo34
- 29 Burn E, Li X, Kostka K, *et al.* Background rates of five thrombosis with thrombocytopenia syndromes of special interest for COVID-19 vaccine safety surveillance: incidence between 2017 and 2019 and patient profiles from 38.6 million people in six European countries. *Pharmacoepidemiol Drug Saf* 2022;31:495–510. 10.1002/pds.5419
- 30 Danwang C, Temgoua MN, Agbor VN, *et al.* Epidemiology of venous thromboembolism in Africa: a systematic review. *J Thromb Haemost* 2017;15:1770–81. 10.1111/jth.13769
- 31 Cohen SL, Gianos E, Barish MA, *et al.* Prevalence and predictors of venous thromboembolism or mortality in hospitalized COVID-19 patients. *Thromb Haemost* 2021;121:1043–53. 10.1055/a-1366-9656
- 32 Cohen CT, Riedl RA, Gowda ST, *et al.* Pulmonary embolism in pediatric and adolescent patients with COVID-19 infection during the SARS-cov-2 delta wave. *Pediatr Blood Cancer* 2022;69:e29721. 10.1002/pbc.29721
- 33 Lee Y, Jehangir Q, Li P, *et al.* Venous thromboembolism in COVID-19 patients and prediction model: a multicenter cohort study. *BMC Infect Dis* 2022;22:462. 10.1186/s12879-022-07421-3
- 34 Bellmont-Montoya S, Riera C, Gil D, *et al.* COVID-19 infection in critically ill patients carries a high risk of venous thromboembolism. *Eur J Vasc Endovasc Surg* 2021;61:628–34. 10.1016/j.ejvs.2020.12.015
- 35 Tan BK, Mainbourg S, Friggeri A, *et al.* Arterial and venous thromboembolism in COVID-19: a study-level meta-analysis. *Thorax* 2021;76:970–9. 10.1136/thoraxjnl-2020-215383
- 36 Jiménez D, García-Sánchez A, Rali P, *et al.* Incidence of vte and bleeding among hospitalized patients with coronavirus disease 2019: a systematic review and meta-analysis. *Chest* 2021;159:1182–96. 10.1016/j.chest.2020.11.005
- 37 Akinyemi RO, Ovbiagele B, Adeniji OA, *et al.* Stroke in Africa: profile, progress, prospects and priorities. *Nat Rev Neurol* 2021;17:634–56. 10.1038/s41582-021-00542-4
- 38 WHO international- News Room feature stories. The Janssen Ad26. COV2.S COVID-19 vaccine: What you need to know. 2022. Available: www.who.int/news-room/feature-stories/detail/the-j-j-covid-19-vaccine-what-you-need-to-know
- 39 Nannoni S, de Groot R, Bell S, *et al.* Stroke in COVID-19: a systematic review and meta-analysis. *Int J Stroke* 2021;16:137–49. 10.1177/1747493020972922
- 40 Bekker L-G, Garrett N, Goga A, *et al.* Effectiveness of the Ad26. COV2.S vaccine in health-care workers in South Africa (the Sisonke study): results from a single-arm, open-label, phase 3B, implementation study. *Lancet* 2022;399:1141–53. 10.1016/S0140-6736(22)00007-1

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjmed-2022-000302>).

Table 1S | Definitions of Vaccine-induced Immune Thrombocytopenia and Thrombosis (VITT) according to Expert Haematology Panel ¹⁷

Definite VITT	All five criteria: <ol style="list-style-type: none"> 1. Onset of symptoms 5–30 days after vaccination against SARS-CoV-2 (or ≤ 42 days in patients with isolated deep-vein thrombosis or pulmonary embolism) 2. Presence of thrombosis 3. Thrombocytopenia (platelet count $< 150 \times 10^9/L$) 4. D-dimer > 4000 FEU 5. Positive anti-PF4 antibodies on ELISA
Probable VITT	D-dimer level > 4000 FEU but one criterion not met (timing, thrombosis, thrombocytopenia, or anti-PF4 antibodies) or d-dimer level unknown or 2000 – 4000 FEU and all other criteria met
Possible VITT	D-dimer level unknown or 2000 – 4000 FEU with one other criterion not met, or two other criteria not met (timing, thrombosis, thrombocytopenia, or anti-PF4 antibodies)
Unlikely VITT	Platelet count $< 150 \times 10^9/L$ and d-dimer level < 2000 FEU, or thrombosis with platelet count $> 150 \times 10^9/L$ and d-dimer level < 2000 FEU, regardless of anti-PF4 antibody result, and alternative diagnosis more likely
Key: ELISA, enzyme-linked immunosorbent assay; FEU, fibrinogen-equivalent unit; PF4, platelet factor 4 and SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.	

Table 2S |Brighton Collaboration interim case definition of Thrombosis with Thrombocytopenia

Confirmed TTS	Presence of major criteria: <ol style="list-style-type: none"> 1. presence of new onset severe thrombocytopenia defined as a platelet count $<50 \times 10^9/L$, and 2. new thrombosis at uncommon site: cerebral veins, splanchnic veins or multiple organ 3. Positive anti-platelet factor 4 antibodies
Probable TTS	Presence of major and minor criteria: <ol style="list-style-type: none"> 1. presence of new onset severe thrombocytopenia defined as a platelet count $<50 \times 10^9/L$, and 2. new thrombosis at common site: pulmonary arteries/veins, limb veins, coronary arteries, cerebral arteries or other arteries/vein OR <ol style="list-style-type: none"> 1. presence of new onset mild to moderate thrombocytopenia defined as a platelet count $<150 \times 10^9/L$, and 2. new thrombosis at uncommon site.
Possible TTS	Presence of minor criteria: <ol style="list-style-type: none"> 1. presence of new onset mild to moderate thrombocytopenia defined as a platelet count $<150 \times 10^9/L$, and 2. new thrombosis at common site. or <ol style="list-style-type: none"> 1. presence of new onset thrombocytopenia defined as a platelet count <150 or $>50 \times 10^9/L$, or $>50\%$ decrease from baseline platelet count and 2. suggestive thrombosis by supporting imaging or laboratory findings or by clinical syndromes in any location 3. D-dimer > 4000 FEU

Supplementary material

Clinical twin study

Data & Methodology

We assessed the risk of 179 conditions post vaccination with AD26.COV2.s amongst Discovery Health clients from 17 February 2021 to 15 September 2021, 18 years and older that had at least 6 months of exposure on Discovery Health. The list of conditions includes 25 conditions analysed in a real world safety study of BNT162b2 vaccination in Israel (Noam Barda, 2021), and a further 154 conditions which are considered immune mediated conditions (Table 4a). Real world safety studies amongst large populations are more likely to provide sufficient volumes to power risk ratio estimates of rare adverse events amongst the vaccinated population compared to randomized control studies which typically have lower volumes. The safety and immunogenicity study of the AD26.COV2.s vaccine was based on 43, 783 participants, split equally between those vaccinated with and those vaccinated with placebo (Sadoff, 2021). Comparison against unvaccinated controls provides greater certainty that identified adverse events are vaccine induced as it provides a risk relative to the population's baseline risk over a consistent timeframe. Registries of voluntary vaccine safety adverse events such as VAERS and the reporting system managed by the South African Health Products Regulatory Agency, unfortunately lack a relevant temporal comparison of underlying adverse events in the unvaccinated general population. This study compares adverse events amongst the vaccinated population against 'clinical twins' who did not receive vaccination. The 'clinical twin' approach was based on the following matching criteria:

- **Age:** clinical twins were matched within one year of age of each other
- **Member's sex:** Male (M) or Female (F)
- **Location:** Hospital service area in South Africa
- **Number of chronic conditions:** 0, 1, 2, 3, 4 or more
- **Number of prior flu vaccinations:** 0, 1, 2, 3, 4, 5 or more. People who had a flu vaccination in the years leading up to and during the COVID-19 pandemic were likely to be more aware of promoting their general health and wellbeing and adhere to preventive measures.
- **One of five medical scheme product options:** Looking at out-of-hospital benefits we divided the 24 plans available through both Discovery Health Medical Scheme and 18 of the Discovery Health administered schemes, into 5 groups.

We assessed the rate of 179 adverse events in the vaccinated population, from the date of vaccination until 42 days after receipt of vaccination to matched unvaccinated ‘clinical twin’ controls. For each condition assessed, clients with a history of the condition were excluded from the analysis i.e. both the vaccinated and matched unvaccinated controls had no prior history of the condition, resulting in the new onset of the condition of interest being analysed after vaccination, relative to the matched baseline population. Individuals with a history of any prior COVID19 vaccination were excluded.

To obtain confidence intervals around risk ratio estimates and the number of adverse events in the study and comparator populations, we bootstrapped amongst our vaccinated population with replacement 500 times. Within each bootstrap vaccinated individuals were matched to a clinical twin, randomly. Once a clinical twin was selected, it was not eligible for selection again within a particular bootstrap. The 95% confidence interval was obtained by the following formula for each of the respective measures analysed (i.e. the risk ratio and number adverse events) : $95\%CI = \mu \pm 1.96\sigma$, where μ is the mean and σ is the standard deviation across bootstraps.

Results

103,315 vaccinated individuals were included in the study. There was a higher proportion of females included (67.6%), as the majority of vaccine recipients were healthcare workers vaccinated under the Sisonke clinical trial, and the nursing profession in South Africa is disproportionately weighted towards females (Table 4b). None of the 179 conditions evaluated amongst the vaccinated population had a statistically significant higher incidence risk ratio relative to unvaccinated controls (Table 4c).

Table 4a: List of analysed conditions

Pulmonary embolism	Evans syndrome	Sympathetic ophthalmia (SO)	Graves disease	Mucha-Habermann disease	Interstitial cystitis (IC)
Kawasaki disease	Immune thrombocytopenic purpura (ITP)	Optic neuritis	Ocular cicatricial pemphigoid	Pernicious anemia (PA)	Autoimmune retinopathy
Acute kidney injury	Thrombocytopenic purpura (TTP)	Leukocytoclastic vasculitis	Takayasu arteritis	Antiphospholipid syndrome	Sperm & testicular autoimmunity
Seizure	Herpes zoster infection	Hypersensitivity pneumonitis	Paraneoplastic cerebellar degeneration (PCD)	Neuromyelitis optica	Herpes simplex infection
Arrhythmia	PANDAS	Lichen sclerosus	Bells palsy	Myelin Oligodendrocyte Glycoprotein Antibody Disorder	Fibromyalgia
Juvenile diabetes (Type 1 diabetes)	Coxsackie myocarditis	Myasthenia gravis	Devic's disease (neuromyelitis optica)	Anti-GBM/Anti-TBM nephritis	Autoimmune oophoritis
Type 1 diabetes	Chronic inflammatory demyelinating polyneuropathy (CIDP)	Achalasia	Lambert-Eaton syndrome	POEMS syndrome	Ligneous conjunctivitis
Anemia	Reactive Arthritis	Mixed connective tissue disease (MCTD)	Dressler's syndrome	Chagas disease	Rheumatoid arthritis
Syncope	Henoch-Schonlein purpura (HSP)	Undifferentiated connective tissue disease (UCTD)	Hypogammaglobulinemia	Lyme disease chronic	Progesterone dermatitis
Glomerulonephritis	Lupus	Erythema nodosum	Polyglandular syndromes type I, II, III	Adult Still's disease	Arthritis or arthropathy
Cerebrovascular accident	Polymyositis	Progressive systemic sclerosis	Churg-Strauss Syndrome (CSS) or Eosinophilic Granulomatosis (EGPA)	Polyarteritis nodosa	Autoimmune orchitis
Deep-vein thrombosis	Subacute bacterial endocarditis (SBE)	Ankylosing spondylitis	Granulomatosis with Polyangiitis	Lichen planus	Hyperthyroidism
Pericarditis	Autoimmune pancreatitis	Alopecia areata	Juvenile myositis (JM)	Scleroderma	Intracranial hemorrhage
Myocardial infarction	Autoimmune encephalomyelitis	Pemphigus	Eosinophilic fasciitis	Vitiligo	Castleman disease (CD)

Vertigo	Multiple sclerosis	Linear IgA disease (LAD)	Essential mixed cryoglobulinemia	Other Immune responses	Lymphopenia
		Goodpastures			Bullous
Myocarditis	Transverse myelitis	syndrome	Pure red cell aplasia (PRCA)	Dermatomyositis pemphigoid	Giant cell Primary sclerosing
IgG4-related	Hidradenitis Suppurativa	myocarditis cholangitis	sclerosing disease	CREST syndrome (HS) (Acne Inversa)	Crohns disease
Fibrosing alveolitis	Polymyalgia rheumatica	Behcets disease	Postmyocardial infarction syndrome	Microscopic polyangiitis (MPA)	Sjögrens syndrome
Autoimmune urticarial	Pars planitis (peripheral uveitis)	Benign mucosal pemphigoid	Inclusion body myositis (IBM)	Autoimmune hepatitis	Addisons disease
Perivenous encephalomyelitis	Uveitis	Other thrombosis	Palindromic rheumatism (PR)	Juvenile arthritis	Retroperitoneal fibrosis
Paresthesia	Vogt-Koyanagi-Harada Disease	Baló disease	Chronic recurrent multifocal osteomyelitis (CRMO)	Motor neuron disease	Autoimmune angioedema
Rheumatic fever	Peripheral neuropathy	Cogans syndrome	Paroxysmal nocturnal hemoglobinuria (PNH)	Dermatitis herpetiformis	Parry Romberg syndrome
Myositis	Discoid lupus	Susacs syndrome	Temporal arteritis/Giant cell arteritis	Neutropenia	Thyroid eye disease (TED)
Lymphadenopathy	IgA Nephropathy	Congenital heart block	Giant cell arteritis (temporal arteritis)	Autoimmune dysautonomia	Sarcoidosis
Axonal & neuronal neuropathy (AMAN)	Primary Biliary Cholangitis	Hashimotos thyroiditis	Amyloidosis	Celiac disease	Scleritis
Thrombocytopenia	Endometriosis	Menieres disease	Agammaglobulinemia	Psoriatic arthritis	Ulcerative colitis (UC)
Guillain-Barre syndrome	Vasculitis	Cicatricial pemphigoid	Restless legs syndrome (RLS)/Stiff person syndrome (SPS)	Relapsing polychondritis	Cold agglutinin disease
Autoimmune myocarditis	Postpericardiotomy syndrome	Moorens ulcer	Herpes gestationis or pemphigoid gestationis (PG)	Psoriasis	Schmidt syndrome
		Raynauds phenomenon			Reflex sympathetic dystrophy
		Multifocal Motor Neuropathy (MMN) or	Neonatal Lupus	Autoimmune inner ear disease (AIED)	

Eosinophilic
esophagitis (EoE)

Parsonage-Turner
syndrome

MMNCB

Appendicitis

Hemolytic anemia

Pyoderma gangrenosum

Tolosa-Hunt syndrome
(THS)

Narcolepsy

Table 4b: Data characteristics

Characteristics	Vaccination analysis	
	Vaccinated Group (N = 103,315)	Control Group (N = 103,315)
Median age (IQR) - yr		
Age group - no. (%)		
<18	(0.0)	71 (0.0)
18-39	55,363 (0.5)	56,803 (0.5)
40-49	26,111 (0.3)	25,533 (0.2)
50-59	14,612 (0.1)	14,130 (0.1)
60-69	5,762 (0.1)	5,430 (0.1)
70-79	1,302 (0.0)	1,199 (0.0)
>=80	166 (0.0)	151 (0.0)
Sex - no (%)		
Female	69,808 (67.6)	69,808 (67.6)
Male	33,507 (32.4)	33,507 (32.4)
Chronic condition count		
0	75,807 (73.4)	75,807 (73.4)
1	17,559 (17.0)	17,559 (17.0)
2	5,482 (5.3)	5,482 (5.3)
3	2,492 (2.4)	2,492 (2.4)
4+	1,977 (1.9)	1,977 (1.9)
No. of influenza vaccinations during preceding 5 yr - no(%)		
0	89,530 (86.7)	89,530 (86.7)
1	10,198 (9.9)	10,198 (9.9)
2	1,992 (1.9)	1,992 (1.9)
3	1,040 (1.0)	1,040 (1.0)
4	479 (0.5)	479 (0.5)
5	77 (0.1)	77 (0.1)

Table 4c: Conditions studied

Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Achalasia	3,227,809	3,227,799	0.0 (0.0 to 0.0)	0.37 (0.0 to 2.0)	0.0 (0.0 to 0.0)

Acute kidney injury	3,189,599	3,189,597	6.32 (2.0 to 12.0)	7.01 (2.0 to 13.0)	1.13 (0.29 to 3.85)
Addisons disease	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)

Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Adult Still's disease	3,231,550	3,231,549	0.0 (0.0 to 0.0)	0.03 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Agammaglobulinemia	3,227,528	3,227,527	0.0 (0.0 to 0.0)	0.04 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Alopecia areata	3,187,925	3,187,813	2.8 (0.0 to 6.0)	7.61 (3.0 to 13.0)	0.42 (0.0 to 1.25)
Amyloidosis	3,232,258	3,232,258	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)
Anemia	2,667,434	2,667,252	37.17 (26.0 to 48.0)	47.8 (35.0 to 61.0)	0.79 (0.51 to 1.16)
Ankylosing spondylitis	3,216,146	3,216,117	0.9 (0.0 to 3.0)	1.58 (0.0 to 4.0)	0.59 (0.0 to 2.0)
Anti-GBM/Anti-TBM nephritis	3,224,889	3,224,886	0.0 (0.0 to 0.0)	0.17 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Antiphospholipid syndrome	3,230,575	3,230,569	0.2 (0.0 to 1.0)	0.36 (0.0 to 2.0)	0.16 (0.0 to 1.0)
Appendicitis	3,070,563	3,070,436	12.34 (6.0 to 21.0)	16.88 (9.0 to 25.0)	0.78 (0.31 to 1.5)
Arrhythmia	3,036,318	3,036,451	23.31 (14.0 to 33.0)	24.79 (16.0 to 35.0)	0.98 (0.52 to 1.71)
Arthritis or arthropathy	2,211,811	2,211,869	66.76 (51.0 to 83.0)	63.92 (50.0 to 79.52)	1.06 (0.74 to 1.49)
Autoimmune angioedema	3,231,821	3,231,821	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Autoimmune dysautonomia	3,228,278	3,228,275	0.73 (0.0 to 2.0)	0.95 (0.0 to 3.0)	0.53 (0.0 to 2.0)
Autoimmune encephalomyelitis	3,224,649	3,224,625	0.0 (0.0 to 0.0)	0.71 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Autoimmune hepatitis	3,230,609	3,230,605	0.0 (0.0 to 0.0)	0.13 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Autoimmune inner ear disease (AIED)	3,202,773	3,202,796	4.38 (1.0 to 9.0)	2.69 (0.0 to 6.0)	1.99 (0.25 to 6.0)
Autoimmune myocarditis	3,225,825	3,225,820	0.0 (0.0 to 0.0)	0.35 (0.0 to 2.0)	0.0 (0.0 to 0.0)

Autoimmune oophoritis	3,147,456	3,147,459	7.47 (3.0 to 13.0)	7.31 (2.0 to 13.0)	1.23 (0.36 to 3.42)
Autoimmune orchitis	3,172,859	3,172,955	8.21 (3.0 to 14.0)	5.35 (1.0 to 10.52)	1.97 (0.43 to 6.0)
Autoimmune pancreatitis	3,219,995	3,219,983	0.0 (0.0 to 0.0)	0.75 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Autoimmune retinopathy	3,145,920	3,145,928	10.34 (4.0 to 17.0)	10.79 (5.0 to 18.0)	1.07 (0.39 to 2.5)
Autoimmune urticarial	2,946,721	2,946,549	19.25 (11.0 to 28.52)	28.56 (19.0 to 39.0)	0.7 (0.34 to 1.12)
Axonal & neuronal neuropathy (AMAN)	3,219,677	3,219,645	0.9 (0.0 to 3.0)	2.75 (0.0 to 6.0)	0.42 (0.0 to 2.0)
Baló disease	3,232,634	3,232,634	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Behcets disease	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Bells palsy	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)

Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Benign mucosal pemphigoid	3,232,258	3,232,258	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Bullous pemphigoid	3,231,658	3,231,656	0.0 (0.0 to 0.0)	0.06 (0.0 to 1.0)	0.0 (0.0 to 0.0)
CREST syndrome	3,231,835	3,231,835	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Castleman disease (CD)	3,225,441	3,225,424	0.0 (0.0 to 0.0)	0.62 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Celiac disease	3,225,081	3,225,068	0.0 (0.0 to 0.0)	0.61 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Cerebrovascular accident	3,147,313	3,147,296	7.6 (3.0 to 13.0)	9.08 (4.0 to 15.52)	0.95 (0.31 to 2.59)
Chagas disease	3,232,180	3,232,180	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Chronic inflammatory demyelinating polyneuropathy (CIDP)	3,225,249	3,225,209	0.0 (0.0 to 0.0)	1.99 (0.0 to 5.0)	0.0 (0.0 to 0.0)
Chronic recurrent multifocal osteomyelitis (CRMO)	3,232,038	3,232,036	0.0 (0.0 to 0.0)	0.12 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Churg-Strauss Syndrome (CSS) or Eosinophilic Granulomatosis (EGPA)	3,232,372	3,232,370	0.0 (0.0 to 0.0)	0.08 (0.0 to 1.0)	0.0 (0.0 to 0.0)

Cicatricial pemphigoid	3,232,258	3,232,258	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Cogans syndrome	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Cold agglutinin disease	3,231,446	3,231,443	0.0 (0.0 to 0.0)	0.09 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Congenital heart block	3,232,134	3,232,134	0.0 (0.0 to 0.0)	0.02 (0.0 to 0.0)	0.0 (0.0 to 0.0)
Coxsackie myocarditis	3,225,824	3,225,824	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Crohns disease	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Deep-vein thrombosis	3,105,790	3,105,901	17.26 (10.0 to 26.0)	12.11 (6.0 to 19.0)	1.56 (0.7 to 3.13)
Dermatitis herpetiformis	3,229,167	3,229,164	0.0 (0.0 to 0.0)	0.16 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Dermatomyositis	3,230,704	3,230,703	0.0 (0.0 to 0.0)	0.04 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Devics disease (neuromyelitis optica)	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Discoid lupus	3,226,370	3,226,390	1.39 (0.0 to 4.0)	0.41 (0.0 to 2.0)	1.39 (0.0 to 4.0)
Dresslers syndrome	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Endometriosis	2,905,162	2,905,088	22.67 (14.0 to 33.0)	22.19 (14.0 to 31.0)	1.07 (0.59 to 1.78)
Eosinophilic esophagitis (EoE)	3,135,425	3,135,350	3.39 (1.0 to 7.0)	6.04 (2.0 to 11.0)	0.68 (0.07 to 2.0)
Eosinophilic fasciitis	3,232,380	3,232,379	0.0 (0.0 to 0.0)	0.05 (0.0 to 1.0)	0.0 (0.0 to 0.0)

Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Erythema nodosum	3,224,679	3,224,701	0.89 (0.0 to 3.0)	0.44 (0.0 to 2.0)	0.84 (0.0 to 3.0)
Essential mixed cryoglobulinemia	3,232,565	3,232,565	0.0 (0.0 to 0.0)	0.01 (0.0 to 0.0)	0.0 (0.0 to 0.0)
Evans syndrome	3,226,511	3,226,534	1.63 (0.0 to 4.0)	0.11 (0.0 to 1.0)	1.62 (0.0 to 3.88)
Fibromyalgia	3,184,600	3,184,625	9.93 (5.0 to 16.0)	10.17 (4.48 to 17.0)	1.11 (0.42 to 2.6)

Fibrosing alveolitis	3,222,706	3,222,690	0.0 (0.0 to 0.0)	0.64 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Giant cell arteritis (temporal arteritis)	3,231,663	3,231,662	0.0 (0.0 to 0.0)	0.12 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Giant cell myocarditis	3,221,587	3,221,573	0.0 (0.0 to 0.0)	1.01 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Glomerulonephritis	3,009,371	3,009,279	14.5 (8.0 to 22.0)	16.95 (9.0 to 25.0)	0.91 (0.41 to 1.71)
Goodpastures syndrome	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Granulomatosis with Polyangiitis	3,232,700	3,232,700	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Graves disease	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Guillain-Barre syndrome	3,226,331	3,226,334	0.92 (0.0 to 3.0)	1.05 (0.0 to 3.0)	0.7 (0.0 to 3.0)
Hashimotos thyroiditis	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Hemolytic anemia	3,231,446	3,231,443	0.0 (0.0 to 0.0)	0.09 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Henoch-Schonlein purpura (HSP)	3,224,358	3,224,337	0.0 (0.0 to 0.0)	0.86 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Herpes gestationis or pemphigoid gestationis (PG)	3,230,789	3,230,784	0.0 (0.0 to 0.0)	0.18 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Herpes simplex infection	2,971,196	2,971,120	17.93 (10.48 to 26.0)	24.76 (16.0 to 36.0)	0.76 (0.36 to 1.29)
Herpes zoster infection	3,076,109	3,076,207	20.08 (11.48 to 30.0)	21.04 (13.0 to 30.0)	0.99 (0.5 to 1.64)
Hidradenitis Suppurativa (HS) (Acne Inversa)	3,208,853	3,208,808	2.15 (0.0 to 5.0)	3.38 (0.0 to 7.0)	0.84 (0.0 to 3.0)
Hypersensitivity pneumonitis	3,214,211	3,214,192	0.0 (0.0 to 0.0)	1.0 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Hyperthyroidism	3,154,251	3,154,146	2.18 (0.0 to 5.52)	5.81 (1.0 to 11.0)	0.5 (0.0 to 2.0)
Hypogammaglobulinemia	3,232,248	3,232,248	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
IgA Nephropathy	3,199,927	3,199,886	0.7 (0.0 to 3.0)	2.24 (0.0 to 6.0)	0.38 (0.0 to 2.0)
IgG4-related sclerosing disease	3,232,137	3,232,136	0.0 (0.0 to 0.0)	0.04 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Immune thrombocytopenic purpura (ITP)	3,227,052	3,227,075	1.63 (0.0 to 4.0)	0.11 (0.0 to 1.0)	1.62 (0.0 to 3.88)
Inclusion body myositis (IBM)	3,231,478	3,231,476	0.0 (0.0 to 0.0)	0.08 (0.0 to 1.0)	0.0 (0.0 to 0.0)

Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Interstitial cystitis (IC)	3,207,419	3,207,450	3.95 (0.0 to 8.0)	2.18 (0.0 to 5.0)	2.24 (0.0 to 7.0)
Intracranial hemorrhage	3,201,232	3,201,238	2.25 (0.0 to 6.0)	2.15 (0.0 to 5.0)	1.24 (0.0 to 4.0)
Juvenile arthritis	3,226,829	3,226,826	0.0 (0.0 to 0.0)	0.18 (0.0 to 1.52)	0.0 (0.0 to 0.0)
Juvenile diabetes (Type 1 diabetes)	3,103,501	3,103,376	5.09 (1.0 to 9.0)	9.33 (4.0 to 15.0)	0.63 (0.13 to 1.75)
Juvenile myositis (JM)	3,232,071	3,232,071	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)
Kawasaki disease	3,232,398	3,232,390	0.0 (0.0 to 0.0)	1.1 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Lambert-Eaton syndrome	3,232,609	3,232,609	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Leukocytoclastic vasculitis	3,209,970	3,209,957	0.52 (0.0 to 2.0)	1.52 (0.0 to 5.0)	0.33 (0.0 to 1.95)
Lichen planus	3,213,599	3,213,626	1.85 (0.0 to 5.0)	1.31 (0.0 to 4.0)	1.39 (0.0 to 4.0)
Lichen sclerosus	3,225,397	3,225,369	0.0 (0.0 to 0.0)	0.96 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Ligneous conjunctivitis	2,332,156	2,332,213	53.94 (40.0 to 69.0)	48.6 (36.0 to 62.52)	1.13 (0.76 to 1.65)
Linear IgA disease (LAD)	3,227,807	3,227,825	0.84 (0.0 to 3.0)	0.32 (0.0 to 2.0)	0.75 (0.0 to 2.0)
Lupus	3,194,102	3,194,050	4.22 (1.0 to 8.0)	5.7 (1.0 to 11.0)	0.96 (0.1 to 3.5)
Lyme disease chronic	3,232,010	3,232,031	0.9 (0.0 to 3.0)	0.2 (0.0 to 1.0)	0.87 (0.0 to 3.0)
Lymphadenopathy	2,884,700	2,884,641	19.04 (11.0 to 27.52)	19.81 (12.0 to 29.0)	1.01 (0.51 to 1.76)
Lymphopenia	3,225,715	3,225,712	0.76 (0.0 to 3.0)	0.39 (0.0 to 2.0)	0.65 (0.0 to 3.0)
Menieres disease	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Microscopic polyangiitis (MPA)	3,228,638	3,228,667	1.05 (0.0 to 3.0)	0.25 (0.0 to 2.0)	0.95 (0.0 to 3.0)
Mixed connective tissue disease (MCTD)	3,229,197	3,229,219	0.94 (0.0 to 3.0)	0.44 (0.0 to 2.0)	0.86 (0.0 to 3.0)
Moorens ulcer	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)

Motor neuron disease	3,229,370	3,229,368	0.0 (0.0 to 0.0)	0.11 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Mucha-Habermann disease	3,232,097	3,232,097	0.0 (0.0 to 0.0)	0.02 (0.0 to 0.0)	0.0 (0.0 to 0.0)
Multifocal Motor Neuropathy (MMN) or MMNCB	3,232,351	3,232,351	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Multiple sclerosis	3,207,647	3,207,615	0.0 (0.0 to 0.0)	1.77 (0.0 to 5.0)	0.0 (0.0 to 0.0)
Myasthenia gravis	3,227,268	3,227,263	0.0 (0.0 to 0.0)	0.37 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Myelin Oligodendrocyte Glycoprotein Antibody Disorder	3,231,805	3,231,795	0.0 (0.0 to 0.0)	0.42 (0.0 to 2.0)	0.0 (0.0 to 0.0)

Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Myocardial infarction	3,192,589	3,192,671	11.59 (6.0 to 19.0)	9.5 (4.0 to 16.0)	1.4 (0.5 to 3.13)
Myocarditis	3,214,286	3,214,271	0.0 (0.0 to 0.0)	1.02 (0.0 to 3.0)	0.0 (0.0 to 0.0)
Myositis	3,103,371	3,103,553	19.04 (12.0 to 27.52)	11.34 (5.0 to 18.0)	1.87 (0.84 to 4.0)
Narcolepsy	3,228,216	3,228,210	0.0 (0.0 to 0.0)	0.25 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Neonatal Lupus	3,225,520	3,225,515	0.0 (0.0 to 0.0)	0.37 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Neuromyelitis optica	3,231,805	3,231,795	0.0 (0.0 to 0.0)	0.42 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Neutropenia	3,224,034	3,224,029	0.0 (0.0 to 0.0)	0.37 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Ocular cicatricial pemphigoid	3,232,154	3,232,148	0.0 (0.0 to 0.0)	0.17 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Optic neuritis	3,218,592	3,218,592	2.0 (0.0 to 5.0)	1.2 (0.0 to 4.0)	1.38 (0.0 to 4.0)
Other Immune responses	3,231,272	3,231,639	11.23 (5.0 to 18.0)	0.86 (0.0 to 3.0)	8.96 (3.0 to 18.0)
Other thrombosis	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
POEMS syndrome	3,231,495	3,231,493	0.0 (0.0 to 0.0)	0.08 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Palindromic rheumatism (PR)	3,229,993	3,229,992	0.0 (0.0 to 0.0)	0.08 (0.0 to 1.0)	0.0 (0.0 to 0.0)

Paraneoplastic cerebellar degeneration (PCD)	3,231,739	3,231,739	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Paresthesia	3,189,133	3,189,283	7.4 (2.48 to 13.0)	2.59 (0.0 to 6.0)	3.48 (0.6 to 10.0)
Paroxysmal nocturnal hemoglobinuria (PNH)	3,232,627	3,232,627	0.0 (0.0 to 0.0)	0.01 (0.0 to 0.0)	0.0 (0.0 to 0.0)
Parry Romberg syndrome	3,230,999	3,230,997	0.0 (0.0 to 0.0)	0.07 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Pars planitis (peripheral uveitis)	3,192,728	3,192,778	4.86 (1.0 to 10.0)	3.18 (0.0 to 7.0)	1.99 (0.25 to 7.0)
Parsonage-Turner syndrome	3,231,637	3,231,637	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Pemphigus	3,229,298	3,229,296	0.0 (0.0 to 0.0)	0.17 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Pericarditis	3,213,907	3,213,906	1.62 (0.0 to 4.52)	1.23 (0.0 to 4.0)	1.15 (0.0 to 4.0)
Peripheral neuropathy	3,229,219	3,229,234	0.85 (0.0 to 3.0)	0.23 (0.0 to 1.0)	0.81 (0.0 to 3.0)
Perivenous encephalomyelitis	3,188,417	3,188,391	1.98 (0.0 to 5.0)	3.0 (0.0 to 7.0)	0.88 (0.0 to 3.51)
Pernicious anemia (PA)	3,223,758	3,223,755	0.0 (0.0 to 0.0)	0.18 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Polyarteritis nodosa	3,229,454	3,229,485	1.05 (0.0 to 3.0)	0.16 (0.0 to 1.0)	0.99 (0.0 to 3.0)
Polyglandular syndromes type I, II, III	3,232,359	3,232,359	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)

Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Polymyalgia rheumatic	3,214,910	3,214,972	3.04 (0.0 to 7.0)	1.23 (0.0 to 4.0)	2.27 (0.0 to 6.0)
Polymyositis	3,227,556	3,227,610	2.91 (0.0 to 6.52)	0.62 (0.0 to 3.0)	2.55 (0.0 to 6.0)
Postmyocardial infarction syndrome	3,232,396	3,232,393	0.0 (0.0 to 0.0)	0.09 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Postpericardiotomy syndrome	3,232,616	3,232,607	0.0 (0.0 to 0.0)	0.45 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Primary Biliary Cholangitis	3,224,739	3,224,734	2.48 (0.0 to 6.0)	1.46 (0.0 to 4.0)	1.62 (0.0 to 5.0)
Primary sclerosing cholangitis	3,224,394	3,224,388	2.48 (0.0 to 6.0)	1.49 (0.0 to 4.0)	1.61 (0.0 to 5.0)

Progesterone dermatitis	2,340,304	2,340,372	68.9 (54.0 to 85.52)	68.57 (51.0 to 86.0)	1.02 (0.72 to 1.41)
Progressive systemic sclerosis	3,228,477	3,228,476	0.0 (0.0 to 0.0)	0.07 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Psoriasis	3,150,517	3,150,499	6.34 (2.0 to 12.0)	7.87 (3.0 to 13.0)	0.96 (0.23 to 2.64)
Psoriatic arthritis	3,150,025	3,150,007	6.34 (2.0 to 12.0)	7.87 (3.0 to 13.0)	0.96 (0.23 to 2.64)
Pulmonary embolism	3,109,556	3,109,669	12.86 (7.0 to 21.0)	10.49 (5.0 to 18.0)	1.37 (0.55 to 3.11)
Pure red cell aplasia (PRCA)	3,231,132	3,231,132	0.0 (0.0 to 0.0)	0.01 (0.0 to 0.0)	0.0 (0.0 to 0.0)
Pyoderma gangrenosum	3,231,641	3,231,641	0.0 (0.0 to 0.0)	0.16 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Raynauds phenomenon	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Reactive Arthritis	3,222,973	3,223,023	3.21 (0.0 to 7.0)	1.03 (0.0 to 3.0)	2.42 (0.0 to 6.0)
Reflex sympathetic dystrophy	3,231,171	3,231,183	1.35 (0.0 to 4.0)	0.14 (0.0 to 1.0)	1.23 (0.0 to 3.0)
Relapsing polychondritis	3,230,175	3,230,180	0.53 (0.0 to 2.0)	0.37 (0.0 to 2.0)	0.5 (0.0 to 2.0)
Restless legs syndrome (RLS)/Stiff person syndrome (SPS)	3,222,348	3,222,312	0.0 (0.0 to 0.0)	2.0 (0.0 to 5.0)	0.0 (0.0 to 0.0)
Retroperitoneal fibrosis	3,229,267	3,229,264	0.0 (0.0 to 0.0)	0.19 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Rheumatic fever	3,208,940	3,208,952	1.57 (0.0 to 4.0)	1.67 (0.0 to 4.0)	0.95 (0.0 to 3.45)
Rheumatoid arthritis	3,085,558	3,085,571	11.77 (5.0 to 19.0)	14.25 (7.0 to 22.0)	0.9 (0.34 to 1.86)
Sarcoidosis	3,220,129	3,220,143	1.0 (0.0 to 3.0)	0.61 (0.0 to 2.0)	0.82 (0.0 to 3.0)
Schmidt syndrome	3,232,359	3,232,359	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Scleritis	3,214,650	3,214,630	0.99 (0.0 to 3.0)	1.75 (0.0 to 5.0)	0.61 (0.0 to 3.0)
Scleroderma	3,228,608	3,228,603	0.0 (0.0 to 0.0)	0.17 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Condition	Average study population exposure (in days)	Average clinical twin exposure (in days)	Adverse events for study population (95% CI)	Adverse events for clinical twin population (95% CI)	Risk Ratio (95% CI)
Seizure	2,984,509	2,984,396	19.37 (12.0 to 28.52)	25.25 (16.0 to 35.0)	0.8 (0.44 to 1.37)

Sjögrens syndrome	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Sperm & testicular autoimmunity	3,213,695	3,213,691	0.96 (0.0 to 3.0)	1.41 (0.0 to 4.0)	0.67 (0.0 to 3.0)
Subacute bacterial endocarditis (SBE)	3,228,351	3,228,343	0.0 (0.0 to 0.0)	0.37 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Susacs syndrome	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Sympathetic ophthalmia (SO)	3,231,701	3,231,733	1.02 (0.0 to 3.0)	0.12 (0.0 to 1.0)	1.06 (0.0 to 3.0)
Syncope	2,961,279	2,961,344	23.19 (14.0 to 33.0)	22.75 (14.0 to 32.52)	1.07 (0.57 to 1.88)
Takayasu arteritis	3,232,734	3,232,734	0.0 (0.0 to 0.0)	0.0 (0.0 to 0.0)	0.0 (nan to nan)
Temporal arteritis/Giant cell arteritis	3,231,857	3,231,855	0.0 (0.0 to 0.0)	0.12 (0.0 to 1.0)	0.0 (0.0 to 0.0)
Thrombocytopenia	3,200,216	3,200,232	2.56 (0.0 to 6.0)	1.88 (0.0 to 5.0)	1.52 (0.0 to 5.58)
Thrombocytopenic purpura (TTP)	3,227,052	3,227,075	1.63 (0.0 to 4.0)	0.11 (0.0 to 1.0)	1.62 (0.0 to 3.88)
Thyroid eye disease (TED)	3,204,746	3,204,729	1.67 (0.0 to 5.0)	2.6 (0.0 to 6.0)	0.82 (0.0 to 3.0)
Tolosa-Hunt syndrome (THS)	3,227,277	3,227,266	0.0 (0.0 to 0.0)	0.55 (0.0 to 2.0)	0.0 (0.0 to 0.0)
Transverse myelitis	3,229,760	3,229,797	1.76 (0.0 to 5.0)	0.26 (0.0 to 1.52)	1.68 (0.0 to 5.0)
Type 1 diabetes	3,103,501	3,103,376	5.09 (1.0 to 9.0)	9.33 (4.0 to 15.0)	0.63 (0.13 to 1.75)
Ulcerative colitis (UC)	3,157,352	3,157,336	4.61 (1.0 to 9.0)	4.73 (1.0 to 10.0)	1.31 (0.23 to 5.0)
Undifferentiated connective tissue disease (UCTD)	3,229,197	3,229,219	0.94 (0.0 to 3.0)	0.44 (0.0 to 2.0)	0.86 (0.0 to 3.0)
Uveitis	3,193,094	3,193,144	4.86 (1.0 to 10.0)	3.18 (0.0 to 7.0)	1.99 (0.25 to 7.0)
Vasculitis	3,208,063	3,208,046	0.52 (0.0 to 2.0)	1.65 (0.0 to 5.0)	0.32 (0.0 to 1.0)
Vertigo	2,864,193	2,864,604	51.75 (39.0 to 66.52)	42.37 (30.0 to 55.52)	1.25 (0.83 to 1.91)
Vitiligo	3,218,418	3,218,424	2.05 (0.0 to 5.0)	1.07 (0.0 to 3.0)	1.52 (0.0 to 4.58)
Vogt-Koyanagi-Harada Disease	3,193,094	3,193,144	4.86 (1.0 to 10.0)	3.18 (0.0 to 7.0)	1.99 (0.25 to 7.0)