






Association between SARS-CoV-2 infection and newly diagnosed hypertension during pregnancy: prospective, population based cohort study

Anne K. Örtqvist ^{1,2}, Maria C. Magnus,³ Elisabeth Dahlqvist,¹ Jonas Söderling,¹ Kari Johansson ^{1,4}, Anna Sandström,^{1,4} Siri E. Håberg,³ Olof Stephansson ^{1,4}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjmed-2022-000465>).

¹Clinical Epidemiology Division, Department of Medicine, Solna, Karolinska Institutet, Stockholm, Sweden

²Department of Obstetrics and Gynecology, Visby Hospital, Visby, Sweden

³Centre for Fertility and Health, Norwegian Institute of Public Health, Oslo, Norway

⁴Department of Women's Health, Division of Obstetrics, Karolinska University Hospital, Stockholm, Sweden

Correspondence to: Dr Anne K. Örtqvist, Clinical Epidemiology Division, Department of Medicine, Solna, Karolinska Institutet, SE-171 76 Stockholm, Sweden; anne.ortqvist@ki.se

Cite this as: *BMJMED* 2023;2:e000465. doi:10.1136/bmjmed-2022-000465

Received: 19 December 2022
Accepted: 26 April 2023

ABSTRACT

OBJECTIVE To study the association between SARS-CoV-2 infection and newly diagnosed hypertension during pregnancy.

DESIGN Prospective, population based cohort study.

SETTING All singleton pregnancies after 22 completed gestational weeks registered in the Swedish Pregnancy Register and the Medical Birth Registry of Norway, from 1 March 2020 to 24 May 2022.

PARTICIPANTS 312 456 individuals available for analysis (201 770 in Sweden and 110 686 in Norway), with pregnancies that reached 42 completed gestational weeks by the end of follow-up in the pregnancy registries, excluding individuals with SARS-CoV-2 infection before pregnancy and those with a diagnosis of pre-existing hypertension or onset of hypertension before 20 gestational weeks.

MAIN OUTCOME MEASURES Newly diagnosed hypertension during pregnancy was defined as a composite outcome of a diagnosis of gestational hypertension, pre-eclampsia, HELLP (haemolysis, elevated liver enzymes, low platelets) syndrome, or eclampsia, from gestational week 20 to one week after delivery. The association between SARS-CoV-2

infection and hypertension during pregnancy was investigated with a stratified Cox proportional hazard model, adjusting for maternal age, body mass index, parity, smoking, region of birth, education, income, coexisting medical conditions, previous hypertension during pregnancy, number of healthcare visits during the past year, and vaccination against SARS-CoV-2. Pre-eclampsia was also analysed as a separate outcome.

RESULTS Of 312 456 individuals available for analysis, 8% (n=24 566) had SARS-CoV-2 infection any time during pregnancy, 6% (n=18 051) had a diagnosis of hypertension during pregnancy, and 3% (9899) had pre-eclampsia. SARS-CoV-2 infection during pregnancy was not associated with an increased risk of hypertension during pregnancy (adjusted hazard ratio 0.99, 95% confidence interval 0.93 to 1.04) or pre-eclampsia (0.98, 0.87 to 1.10). The results were similar for SARS-CoV-2 infection in all gestational trimesters and in different time periods that corresponded to dominance of different variants of the SARS-CoV-2 virus.

CONCLUSIONS This population based study did not find any evidence of an association between SARS-CoV-2 infection during pregnancy and an increased risk of hypertension during pregnancy or pre-eclampsia.

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Previous studies have suggested that SARS-CoV-2 infection in pregnancy is a risk factor for newly diagnosed hypertension during pregnancy, including pre-eclampsia
- ⇒ Covid-19 disease is a spectrum of signs and symptoms in pregnant individuals that might resemble pre-eclampsia
- ⇒ Any associations between SARS-CoV-2 infection and hypertension during pregnancy could reflect closer monitoring and a higher probability of being tested for SARS-CoV-2 infection in those with early symptoms of hypertensive disorders in pregnancy

WHAT THIS STUDY ADDS

- ⇒ Although SARS-CoV-2 infection in pregnancy is a risk factor for several adverse pregnancy outcomes, no evidence of an association between SARS-CoV-2 infection and newly diagnosed hypertension during pregnancy was found

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE, OR POLICY

- ⇒ In this study, timing of infection and onset of hypertension during pregnancy was accounted for, which has been lacking in previous studies
- ⇒ As SARS-CoV-2 in pregnancy is related to other adverse outcomes in pregnancy, vaccination of pregnant women is still recommended

Introduction

Newly diagnosed hypertension during pregnancy after 20 gestational weeks includes gestational hypertension and pre-eclampsia, affects about 6-8% of the pregnant population worldwide, and is a leading cause of maternal and perinatal morbidity and mortality.^{1 2} Current knowledge suggests that pregnancy does not increase susceptibility to the SARS-CoV-2 virus, but might worsen the manifestations of covid-19 compared with non-pregnant individuals of the same age.³

Severe covid-19 disease has been shown to have multiorgan effects and symptomatology, similar to those seen in individuals with hypertension during pregnancy. These effects include newly diagnosed hypertension, kidney injury, liver involvement, haematological complications, pathological placentas, and endothelial dysfunction, and therefore a biological link between infection by the SARS-CoV-2 virus and hypertension during pregnancy, specifically pre-eclampsia, has been suggested.⁴⁻⁶ A

recent systematic review and meta-analysis showed that SARS-CoV-2 infection during pregnancy was associated with a significant increase in the odds of developing pre-eclampsia (pooled adjusted odds ratio 1.58, 95% confidence interval 1.39 to 1.80; $P < 0.001$).⁶ The renin-angiotensin system is an important regulator of placental function because it modulates proliferation of trophoblasts, angiogenesis, and blood flow. It has been shown that the SARS-CoV-2 virus binds to the angiotensin converting enzyme 2 receptor expressing maternal and fetal cells in the placenta, leading to changes in the renin-angiotensin system, which might have a role in the suggested association between SARS-CoV-2 infection and pre-eclampsia.⁶

Severe covid-19 disease and pre-eclampsia share the same risk factors: pre-existing hypertension, obesity, advanced maternal age, diabetes,^{7, 8} chronic kidney disease,⁹ and systemic lupus erythematosus.¹⁰ These factors could contribute to unmeasured confounding and thus cause spurious associations. Another potential bias is that pregnancies with early symptoms of hypertension are monitored more closely and have a higher probability of being tested for SARS-CoV-2 than pregnancies without adverse symptoms. This surveillance bias could introduce non-random misclassification of the risk factor and reversed causation. Only one¹¹ of the included studies in the meta-analysis⁶ considered the time between infection and pre-eclampsia (ie, temporality).

Population based healthcare and quality registers from Sweden and Norway have detailed information on dates of positive test results for SARS-CoV-2 infection and confirmed diagnoses of hypertension during pregnancy. In this study, we looked at the association between SARS-CoV-2 infection and subsequent hypertension during pregnancy and pre-eclampsia, taking into account temporality and confounding.

Methods

Study population and design

This prospective population based cohort study included all singleton pregnancies after 22 completed gestational weeks registered in the Swedish Pregnancy Register¹² and the Medical Birth Registry of Norway¹³ from 1 March 2020 to 24 May 2022. Access to free perinatal care is similar in the two countries. With the unique personal identification number assigned to all citizens, individual data on births were linked to registry data to collect information on infection, outcomes, and other population characteristics. Online supplemental file describes the registers in more detail.

To avoid oversampling of short pregnancies towards the end of follow-up (preterm births would be registered as births whereas ongoing pregnancies would not yet be registered if delivery was after the end of follow-up), only pregnancies with the

possibility to reach 42 completed gestational weeks by the end of follow-up in the pregnancy registries were included. Individuals with SARS-CoV-2 infection before pregnancy were excluded to ensure that we looked at SARS-CoV-2 infection during pregnancy with onset of hypertension after 20 weeks of gestation. We also excluded those with a diagnosis of pre-existing hypertension or onset of hypertension before 20 gestational weeks, defined according to the ICD-10 (International Classification of Diseases, 10th revision) or Anatomical Therapeutic Chemical codes listed in the online supplemental file.

SARS-CoV-2 infection

Data on SARS-CoV-2 infection were identified from the Public Health Agency in Sweden¹⁴ and the Norwegian Surveillance System for Communicable Diseases.¹⁵ From February 2020 onwards, reporting of all laboratory confirmed polymerase chain reaction (PCR) test results for SARS-CoV-2 was mandatory in both countries.^{15, 16} In this study, individuals were defined as being infected if they had a positive PCR test result for SARS-CoV-2 between the estimated date of conception and the date of delivery. The exact date of when the PCR test was performed was used in the analysis as a proxy for onset of infection. Individuals with a positive test result for SARS-CoV-2 after the outcome of hypertension during pregnancy were treated as censored. Although limited by the absence of complete variant sequencing data, we identified four time periods when different variants of the SARS-CoV-2 virus were dominant: index (1 March 2020-31 January 2021; the first period of the pandemic with the newly identified virus before variants of the virus were identified), alpha variant (1 February 2021-30 June 2021), delta variant (1 July 2021-31 December 2021), and omicron variant (1 January 2022-24 May 2022).¹⁷ Testing strategies and vaccination policies for pregnant individuals in Sweden and Norway have previously been described^{16, 18} and are summarised in the online supplemental file. In this study, we used information from national vaccination registries¹⁸ on dates and number of doses of vaccine against the SARS-CoV-2 virus.

Hypertension during pregnancy and pre-eclampsia

Information on outcomes was collected from the Swedish Pregnancy Register, Medical Birth Registry of Norway, and national patient registers.^{15, 19} Outcomes were hypertension during pregnancy, defined as having a registered ICD-10 code for gestational hypertension (O13.9), pre-eclampsia (O14.0, O14.1, and O14.9), HELLP (haemolysis, elevated liver enzymes, low platelets) syndrome (O14.2), or eclampsia (O15), from 20 gestational weeks to one week postpartum; and pre-eclampsia, which excluded gestational hypertension, but included HELLP and eclampsia. The first date of the registered ICD code was used in the analysis as the

onset of outcome. Online supplemental file describes the definitions and registries used for the outcomes.

Covariates

Population characteristics were derived from the Swedish Pregnancy Register, Medical Birth Registry of Norway, national patient registries, Statistics Sweden,²⁰ and Statistics Norway.²¹ Characteristics considered potential confounders and subsequently adjusted for were maternal age at the start of pregnancy (continuous), body mass index (continuous), parity (0, 1, and ≥ 2), income (top, middle, and bottom thirds), region of birth (Scandinavia, other European countries, Middle East/Africa, and other, including North America, South America, Latin America, Asia, Australia, and New Zealand), highest educational level (< 9 , 10-12, and > 12 years of schooling), smoking in early pregnancy, living with a partner, pre-existing and gestational diabetes, chronic kidney disease, systemic lupus erythematosus, history of hypertension during pregnancy in previous pregnancies, number of visits to an outpatient healthcare facility within one year of conception, and vaccination against SARS-CoV-2 (vaccinated before conception or vaccinated during pregnancy, treated as time varying). For the Norwegian analysis, information on the number of visits to an outpatient healthcare facility within one year of conception was not available.

Statistical analysis

To investigate the association between (date of) SARS-CoV-2 infection and onset of hypertension during pregnancy and pre-eclampsia (date of diagnosis), we used a Cox proportional hazard model with gestational age as the underlying time scale and stratified by the estimated date of conception. Date of conception and gestational age at birth were estimated from the embryo transfer date, routine ultrasound assessment at about 18 gestational weeks (95%), or by using the date of the last menstrual period if other estimates were missing. The stratified Cox analysis allows for different baseline hazards for each group of pregnant individuals with an estimated date of conception on the same date, thus taking into account variations in baseline risks of SARS-CoV-2 (as well as changes in testing capacity in relation to different calendar time periods) during the course of the study. To deal with immortal time bias (ie, to avoid misclassification of time of infection caused by treating infected during follow-up as being infected during the whole follow-up) SARS-CoV-2 infection was modelled as a time varying exposure. Thus those with SARS-CoV-2 infection after the onset of hypertension during pregnancy were treated as censored. The Cox model also deals with timing of events and ensures that the test date was before the first registration of any hypertension during pregnancy.

We analysed the association between SARS-CoV-2 infection and hypertension during pregnancy, and SARS-CoV-2 infection and pre-eclampsia. In secondary analyses, we investigated SARS-CoV-2 infection by gestational trimester and time period of infection based on the dominant variant of the SARS-CoV-2 virus (index, alpha, delta, or omicron). Also, we differentiated the associations between SARS-CoV-2 infection and the risk of hypertension during pregnancy and pre-eclampsia according to whether individuals were vaccinated or not before infection with the SARS-CoV-2 virus.

Results are reported as crude and adjusted hazard ratios with 95% confidence intervals. The proportional hazards assumption was tested with Schoenfeld's residuals. We did not impute missing information on covariates because individuals with missing information, particularly for education and income, were more likely to be immigrants; limited information on immigrants is available to impute missing values. Instead, missing data were managed with a missing indicator variable in the adjusted models for the variables with missing information.

A meta-analysis, with the Knapp-Hartung random effect model, was conducted to estimate the combined adjusted hazard ratio for the primary analyses in Sweden and Norway. Heterogeneity was evaluated with Q statistics (with P values), τ^2 , and I^2 .²² The Swedish analyses were performed with SAS version 9.4 (SAS institute) and the Norwegian data were analysed with Stata version 15 and R version 4.1.2 (R Core Team, 2021), with the Meta package for the meta-analysis.

Sensitivity analyses

To assess the reliability of the results, sensitivity analyses were performed: firstly, individuals with the outcome of hypertension during pregnancy, with a positive test result for SARS-CoV-2 within seven days of the outcome (before or after) were excluded from the main analysis of the association between SARS-CoV-2 infection and hypertension during pregnancy, to further reduce problems with temporality; secondly, because information on coexisting medical conditions and income were limited after 10 October 2021 in the Swedish data, the analysis was conducted with the end of follow-up as this date. The estimated parameters were then compared with estimated parameters from the main analysis where information on coexisting medical conditions and income were limited between 10 October 2021 and 24 May 2022; and thirdly, HELLP syndrome and eclampsia were excluded from the outcome of pre-eclampsia.

Patient and public involvement

Patients and the public were not personally involved in the research process of this study, except for




Figure 1 | Development of cohorts for study of hypertension during pregnancy according to SARS-CoV-2 infection

contributing to data in the national registries. During the study, a high priority was given to a fast study process in order to obtain results quickly for policy makers and clinicians, which precluded patient and public involvement at this point. The results have been orally presented at national conferences for clinicians, and popular scientific summaries of the findings will also be published on web pages for the pregnancy registries.

Results

The selection of study participants is shown in figure 1. Table 1 presents the characteristics of participants according to SARS-CoV-2 infection during pregnancy, and includes a description of the proportion of missing data. Of the 312 456 individuals available for analysis (201 770 in Sweden and 110 686 in Norway), 24 566 (8%) were infected with the SARS-CoV-2 virus during pregnancy, 15% during the first trimester, 31% in the second trimester, and 53% in the third trimester (online supplemental eTable 1). Among those infected with the SARS-CoV-2 virus during pregnancy, most individuals (n=6655, 35%) in the Swedish cohort had a positive test result during the index period, followed by the omicron (n=5709, 30%), alpha (n=4855, 27%), and delta (n=1744, 9%) periods (online supplemental eTable 2). In Norway, most people (n=3211, 57%)

were infected during the omicron period, followed by the delta (1221, 22%), alpha (n=590, 11%), and index (n=574, 10%) periods.

Overall, independent of SARS-CoV-2 infection, 6% of individuals available for analysis (11 620 in Sweden and 6431 in Norway) were defined as having hypertension during pregnancy, and 3% with pre-eclampsia (5891 in Sweden and 4008 in Norway) (online supplemental eTable 3). Among those infected with the SARS-CoV-2 virus, independent of infection before the outcome, 5% also had hypertension during pregnancy (1029 in Sweden and 309 in Norway) and 3% had pre-eclampsia (501 in Sweden and 186 in Norway). Mean time from SARS-CoV-2 infection any time during pregnancy to first registration of hypertension during pregnancy was 88.2 days (standard deviation 78.5, range -86 to 295) in Sweden and 57.1 days (58.0, -30 to 272) in Norway (online supplemental eTable 4). For pre-eclampsia, mean time was 86.3 days (77.8, -77 to 286) in Sweden and 60.5 days (57.6, -20 to 272) in Norway.

Figure 2 presents crude and adjusted hazard ratios and 95% confidence intervals for hypertension during pregnancy and pre-eclampsia according to SARS-CoV-2 infection in Sweden and Norway, and in Sweden and Norway combined. In the crude and adjusted analyses, we found no increased risk of hypertension during pregnancy (combined adjusted hazard ratio 0.99, 95% confidence interval 0.93 to 1.04; $I^2=0\%$, $Q=0.02$, P value for heterogeneity=0.89; $\tau^2=0$) or pre-eclampsia (0.98, 0.87 to 1.10; $I^2=0\%$, $Q=0.04$, P value for heterogeneity=0.85, $\tau^2=0$), with similar estimates for Sweden and Norway in analyses by country.

Table 2 shows crude and adjusted hazard ratios and 95% confidence intervals for hypertension during pregnancy and pre-eclampsia according to SARS-CoV-2 infection by trimester, time periods corresponding to the dominant variant of the SARS-CoV-2 virus, and vaccination status. We found no evidence for an association between hypertension during pregnancy (adjusted hazard ratio range 0.94-1.03 in Sweden, and 0.65-1.04 in Norway) and individuals infected by SARS-CoV-2 in the first, second, or third trimester. Similarly, we found no increased risk of pre-eclampsia (adjusted hazard ratio range 0.91-1.05 in Sweden and 0.77-1.04 in Norway). Also, we found no difference in the risk of hypertension during pregnancy (adjusted hazard ratio range 0.80-1.08) or pre-eclampsia (0.74-1.13) according to SARS-CoV-2 infection during time periods corresponding to the different dominant variants of the virus, or according to infection before or after vaccination for covid-19 (0.97-1.11).

In the sensitivity analysis excluding individuals (114 in Sweden and 40 in Norway) with SARS-CoV-2 infection within seven days of the outcome of hypertension during pregnancy, the adjusted

Table 1 | Participant characteristics according to SARS-CoV-2 infection during pregnancy, March 2020-May 2022, in Sweden and Norway

Characteristics*	Sweden		Norway	
	No infection (n=182 810)	SARS-CoV-2 infection (n=18 960)	No infection (n=105 090)	SARS-CoV-2 infection (n=5596)
SARS-CoV-2 vaccination:				
Vaccinated before pregnancy	5787 (3.2)	1839 (9.7)	643 (0.6)	406 (7.3)
One dose of vaccine during pregnancy	27 533 (15.1)	4284 (22.6)	15 586 (14.8)	2747 (49.1)
Not vaccinated	149 490 (81.8)	12 837 (67.7)	88 861 (84.6)	2443 (43.7)
Maternal age (years):				
<20	2046 (1.1)	149 (0.8)	902 (0.9)	36 (0.6)
20-24	18 336 (10.0)	1852 (9.8)	10 575 (10.1)	478 (8.5)
25-29	60 887 (33.3)	6396 (33.7)	35 796 (34.1)	1836 (32.8)
30-34	66 754 (36.5)	6980 (36.8)	39 055 (37.2)	2197 (39.3)
35-39	28 993 (15.9)	2972 (15.7)	15 992 (15.2)	899 (16.1)
≥40	5794 (3.2)	611 (3.2)	2770 (2.6)	150 (2.7)
Body mass index:				
<18.5	4083 (2.2)	330 (1.7)	3253 (3.1)	158 (2.8)
18.5-<25	94 922 (51.9)	9345 (49.3)	57 759 (55.0)	3066 (54.8)
25-<30	48 708 (26.6)	5375 (28.3)	23 068 (22.0)	1299 (23.2)
≥30	28 039 (15.3)	3266 (17.2)	13 765 (13.1)	737 (13.2)
Missing	7058 (3.9)	644 (3.4)	7245 (6.9)	336 (6.0)
Parity:				
0	79 398 (43.4)	7496 (39.5)	46 277 (44.0)	2092 (37.4)
1	67 885 (37.1)	7142 (37.7)	38 660 (36.8)	2061 (36.8)
≥2	35 527 (19.4)	4322 (22.8)	20 153 (19.2)	1443 (25.8)
Educational level (years):				
<9	16 812 (9.2)	1465 (7.7)	13 722 (10.1)	905 (16.2)
10-12	66 543 (36.4)	6845 (36.1)	20 138 (19.2)	1095 (9.6)
>12	89 896 (49.2)	9069 (47.8)	61 255 (58.3)	2878 (51.4)
Missing	9559 (5.2)	1581 (8.3)	9975 (9.5)	718 (12.8)
Income:				
Bottom third	46 717 (25.6)	3363 (17.7)	45 898 (43.7)	2758 (49.3)
Middle third	46 548 (25.5)	3531 (18.6)	30 324 (28.9)	1413 (25.3)
Top third	46 859 (25.6)	3220 (17.0)	23 497 (22.4)	1074 (19.2)
Missing	40 387 (22.1)	8700 (45.9)	5371 (5.1)	351 (6.3)
Birth region:				
Scandinavia	126 378 (69.1)	12 755 (67.3)	78 400 (74.6)	3613 (64.6)
Other European countries	14 574 (8.0)	1500 (7.9)	11 728 (11.2)	767 (13.7)
Middle East/Africa	28 826 (15.8)	3205 (16.9)	6987 (6.7)	764 (13.7)
Other	8518 (4.7)	661 (3.5)	7960 (7.6)	451 (8.1)
Missing	4514 (2.5)	839 (4.4)	15 (0.01)	1 (0.02)
Smoking status:				
Non-smoker	170 485 (93.3)	17 725 (93.5)	89 050 (84.7)	4678 (83.6)
Smoker	6101 (3.3)	533 (2.8)	5269 (5.0)	263 (4.7)
Missing	6224 (3.4)	702 (3.7)	10 771 (10.3)	655 (11.7)
Living with partner:				
Yes	165 110 (90.3)	17 337 (91.4)	99 197 (94.4)	5248 (93.8)
No	13 727 (7.5)	1270 (6.7)	4153 (4.0)	258 (4.6)
Missing	3973 (2.2)	353 (1.9)	1740 (1.7)	90 (1.6)
Coexisting medical conditions:				
Gestational diabetes	8184 (4.5)	846 (4.5)	6802 (6.5)	379 (6.8)
Pre-existing diabetes mellitus	1994 (1.1)	229 (1.2)	790 (0.8)	31 (0.6)
Chronic kidney disease	645 (0.4)	66 (0.3)	575 (0.6)	33 (0.6)
Systemic lupus erythematosus	259 (0.1)	30 (0.2)	88 (0.1)	2 (0.04)
History of hypertension during pregnancy:				

Continued

Table 1 Continued

Characteristics*	Sweden		Norway	
	No infection (n=182 810)	SARS-CoV-2 infection (n=18 960)	No infection (n=105 090)	SARS-CoV-2 infection (n=5596)
Yes	1413 (0.8)	112 (0.6)	3517 (3.4)	199 (3.6)
Missing	40 387 (22.1)	8700 (45.9)	0	0
No of visits to outpatient care one year before conception:				
0	87 900 (48.1)	6232 (32.9)	n/a	n/a
1-9	51 839 (28.4)	3859 (20.4)	n/a	n/a
10-29	2637 (1.4)	164 (0.9)	n/a	n/a
≥30	47 (0.0)	5 (0.0)	n/a	n/a
Missing	40 387 (22.1)	8700 (45.9)	105 090 (100.0)	5596 (100.0)
Gestational age at birth:				
Week 22+0 to 36+6	7686 (4.2)	796 (4.2)	4726 (4.5)	195 (3.5)
Week 37+0 to 39+6	83 388 (45.6)	8789 (46.4)	43 972 (41.8)	2415 (43.2)
Week ≥40+0	91 736 (50.2)	9375 (49.4)	56 392 (53.7)	2986 (53.4)

Data are numbers (percentages).

*A missing indicator was used in adjusted analyses for body mass index, education, income, birth region, smoking status, history of hypertension during pregnancy, and visits to outpatient care.

hazard ratio was 0.91 (95% confidence interval 0.85 to 0.98) in Sweden and 0.91 (0.78 to 1.04) in Norway. Comparing estimates from Swedish data with complete data on coexisting medical conditions available up to 11 October 2021 with estimates from data with follow-up to 24 May 2022, we found no differences in the results (results not shown). When individuals with HELLP syndrome and eclampsia (176 in Sweden and 165 in Norway) were excluded from the definition of pre-eclampsia, we found only minor changes in effect estimates (results not shown) and the conclusion remained.

Discussion

Principal findings

Our results, based on population based cohorts from Sweden and Norway of 312 456 individuals giving birth during the first two years of the pandemic,

showed no increased risk of hypertension during pregnancy or pre-eclampsia after infection with the SARS-CoV-2 virus during pregnancy. The results were similar for SARS-CoV-2 infection in any trimester, during times corresponding to the different dominant variants of the SARS-CoV-2 virus, and when adjusting for vaccination status.

Previous studies

A recent systematic review and meta-analysis, including 28 studies,⁶ found a consistent association between SARS-CoV-2 infection and pre-eclampsia in several countries with different study designs. Discrepancies between the results from the overall meta-analysis, the individual studies included, and our study could be attributed to several factors. We controlled for a large number of confounding factors prospectively reported to national registers, thereby




Figure 2 | Meta-analysis of hazard ratios for hypertension during pregnancy and pre-eclampsia according to SARS-CoV-2 infection. Adjusted hazard ratios were adjusted for maternal age, body mass index, parity, income, region of birth, educational level, smoking status, living with a partner, gestational and pre-existing diabetes, chronic kidney disease, systemic lupus erythematosus, history of hypertension during pregnancy, number of visits to an outpatient healthcare facility within one year of conception (Swedish data only), and vaccination against SARS-CoV-2. Hypertension during pregnancy: heterogeneity Q=0.02 (P=0.89), $\tau^2=0$, $I^2=0\%$. Pre-eclampsia: heterogeneity Q=0.04 (P=0.85), $\tau^2=0$, $I^2=0\%$. CI=confidence interval

Table 2 | Hazard ratios for hypertension during pregnancy according to SARS-CoV-2 infection in different trimesters, time periods corresponding to the dominant variant of the virus, and vaccination status in Sweden and Norway

SARS-CoV-2 infection	Patients (No (%))	Follow-up time after infection (days)	Crude hazard ratio (95% CI)	Adjusted hazard ratio* (95% CI)
Sweden: hypertension during pregnancy				
No infection	10 568 (5.8)		Reference	Reference
Positive test result in:				
First trimester	175 (5.4)	347 634	0.98 (0.84 to 1.13)	0.94 (0.81 to 1.10)
Second trimester	364 (5.6)	725 441	0.96 (0.86 to 1.07)	0.97 (0.87 to 1.08)
Third trimester	489 (5.3)	349 640	1.01 (0.91 to 1.12)	1.03 (0.93 to 1.15)
Index variant	429 (6.5)	651 148	1.09 (0.98 to 1.20)	1.08 (0.98 to 1.20)
Alpha variant	244 (5.0)	355 105	0.93 (0.82 to 1.07)	0.91 (0.79 to 1.04)
Delta variant	73 (4.2)	130 816	0.78 (0.61 to 0.99)	0.86 (0.67 to 1.10)
Omicron variant	282 (4.9)	285 646	0.93 (0.81 to 1.07)	0.97 (0.84 to 1.11)
At least one dose of vaccine before infection	261 (5.4)	272 427	1.03 (0.89 to 1.18)	1.01 (0.87 to 1.16)
Not vaccinated	767 (5.4)	1 150 288	0.97 (0.90 to 1.05)	0.98 (0.91 to 1.06)
Sweden: pre-eclampsia				
No infection	5 376 (2.9)		Reference	Reference
Positive test result in:				
First trimester	90 (2.8)	348 969	0.97 (0.78 to 1.20)	0.93 (0.75 to 1.16)
Second trimester	173 (2.7)	728 416	0.91 (0.78 to 1.06)	0.91 (0.78 to 1.06)
Third trimester	237 (2.6)	352 374	1.01 (0.87 to 1.18)	1.05 (0.90 to 1.23)
Index variant	231 (3.5)	654 192	1.15 (1.00 to 1.33)	1.13 (0.98 to 1.31)
Alpha variant	120 (2.5)	356 447	0.88 (0.73 to 1.07)	0.86 (0.71 to 1.05)
Delta variant	33 (1.9)	131 436	0.73 (0.51 to 1.05)	0.79 (0.55 to 1.14)
Omicron variant	116 (2.0)	287 684	0.81 (0.65 to 1.00)	0.85 (0.68 to 1.06)
At least one dose of vaccine before infection	107 (2.2)	274 403	0.90 (0.72 to 1.12)	0.92 (0.74 to 1.14)
Not vaccinated	393 (2.8)	1 155 356	0.98 (0.88 to 1.09)	0.98 (0.88 to 1.10)
Norway: hypertension during pregnancy				
No infection	6082 (5.5)		Reference	Reference
Positive test result in:				
First trimester	18 (4.2)	59 065	0.66 (0.41 to 1.05)	0.65 (0.40 to 1.08)
Second trimester	69 (5.5)	143 498	0.81 (0.63 to 1.03)	0.97 (0.75 to 1.25)
Third trimester	190 (5.3)	144 182	0.90 (0.77 to 1.06)	1.04 (0.89 to 1.23)
Index variant	22 (3.9)	58 534	0.71 (0.47 to 1.09)	0.80 (0.52 to 1.24)
Alpha variant	24 (4.2)	56 803	0.80 (0.53 to 1.20)	0.95 (0.63 to 1.43)
Delta variant	60 (5.2)	97 198	0.81 (0.63 to 1.06)	0.92 (0.71 to 1.21)
Omicron variant	171 (5.8)	134 210	0.92 (0.77 to 1.09)	1.06 (0.89 to 1.27)
At least one dose of vaccine before infection	169 (5.9)	153 918	0.92 (0.77 to 1.09)	1.01 (0.85 to 1.21)
Not vaccinated	108 (4.3)	192 827	0.79 (0.65 to 0.96)	0.97 (0.79 to 1.18)
Norway: pre-eclampsia				
No infection	3817 (3.5)		Reference	Reference
Positive test result in:				
First trimester	13 (3.0)	59 077	0.80 (0.46 to 1.38)	0.77 (0.42 to 1.40)
Second trimester	43 (3.4)	143 792	0.79 (0.58 to 1.08)	0.94 (0.69 to 1.29)
Third trimester	113 (3.1)	144 880	0.90 (0.73 to 1.10)	1.04 (0.84 to 1.28)
Index variant	13 (2.3)	58 677	0.69 (0.40 to 1.20)	0.74 (0.42 to 1.32)
Alpha variant	13 (2.3)	56 825	0.69 (0.40 to 1.20)	0.83 (0.48 to 1.44)
Delta variant	41 (3.5)	97 415	0.89 (0.65 to 1.22)	0.99 (0.71 to 1.38)
Omicron variant	102 (3.4)	134 832	0.92 (0.73 to 1.16)	1.08 (0.85 to 1.36)
At least one dose of vaccine before infection	107 (3.7)	154 563	0.98 (0.76 to 1.22)	1.11 (0.89 to 1.38)
Not vaccinated	62 (2.5)	193 186	0.73 (0.57 to 0.95)	0.87 (0.67 to 1.14)

CI=confidence interval.

*Adjusted for maternal age, body mass index, parity, income, region of birth, educational level, smoking status, living with a partner, gestational and pre-existing diabetes, chronic kidney disease, systemic lupus erythematosus, history of hypertension during pregnancy, number of visits to an outpatient healthcare facility within one year of conception (Swedish data only), and vaccination against SARS-CoV-2.

precluding recall bias and reducing the risks of residual confounding, whereas only about half of the studies in the meta-analysis controlled for potential confounders.

Most previous studies were limited to small study populations, with a restricted total number of participants infected with the SARS-CoV-2 virus.⁶ In the meta-analysis based on adjusted odds ratios, nine of 11 studies were cohort studies, including 8135 participants, where 36% were defined as infected with the SARS-CoV-2 virus. The other two studies,^{23,24} comprising 99% (748 526 participants of whom 1% were defined as infected with the SARS-CoV-2 virus) of the total included participants in the adjusted analysis, used a cross sectional design, making it difficult to assess whether SARS-CoV-2 infection preceded hypertension during pregnancy.

Temporality is fundamental when assessing causality.^{25,26} Only one previous study investigating the association between SARS-CoV-2 and hypertension during pregnancy took account of temporality.¹¹ In that study of 249 participants, early (before 32 gestational weeks; hazard ratio 2.17, 95% confidence interval 1.11 to 4.24), but not late (1.68, 0.79 to 3.57), infection was significantly associated with hypertension during pregnancy). The authors hypothesised that participants with infection closer to term did not have a chance to develop clinical outcomes of hypertension during pregnancy. In line with this hypothesis, whether it is biologically plausible that an individual with a positive test result for SARS-CoV-2 at birth (even if the pregnant individual might have been infected some time before that) had sufficient time to develop hypertension during pregnancy is unclear. In our study, we found no difference in the results independent of whether infection occurred in the first, second, or third trimester, which makes it unlikely that our lack of association was caused by too short a time for the virus to cause hypertension during pregnancy.

Another potential reason for spurious associations is that individuals with early symptoms of hypertension during pregnancy are monitored more closely and thus have a higher probability of being tested for SARS-CoV-2 than those without hypertension during pregnancy. This practice can cause surveillance bias and non-random misclassification of infection, which could induce reversed causation. Also, pregnant individuals with severe covid-19 can develop a pre-eclampsia-like syndrome²⁷ that might be distinguished from pre-eclampsia by ultrasound assessment and specific angiogenic factors,²⁸ but not always from clinical symptomatology. Hence misclassification of outcome could have occurred. In a sensitivity analysis, to further study temporality and the co-occurrence of infection and hypertension during pregnancy, we excluded individuals with hypertension during pregnancy that tested positive for SARS-CoV-2 within seven days of the outcome,

and found no increased risk of hypertension during pregnancy after infection.

Studies have reported that the delta variant of the SARS-CoV-2 virus could be associated with higher rates of maternal mortality²⁹ and severe maternal morbidity than other variants.³⁰ But another study found no significant difference in pre-eclampsia by variant type.³¹ Whole genome sequencing data of the different variants of the SARS-CoV-2 virus were missing at the population level in Sweden and Norway. When we investigated the different time periods corresponding to the dominant variants,¹⁷ however, we found no difference in the results.

Strengths and limitations

A major strength of our study was the large population based cohort of prospectively collected data from two Nordic countries, allowing the analyses to account for the timing between infection and outcome, as well as controlling for potential confounders, with consistent results. We also assessed SARS-CoV-2 infection in different trimesters and included national data on vaccination against SARS-CoV-2 in our analysis. This approach is important as the vaccine will influence the probability and severity of SARS-CoV-2 infection.

This study had some limitations. Firstly, misclassification because of unidentified patients with SARS-CoV-2 infection could be a problem with this study design based on registry data. We used population based data on positive test results for SARS-CoV-2 infection verified by PCR as the risk factor, but testing capacity changed over time, and only those with severe symptoms of covid-19 were tested initially.¹⁶ Between June 2020 and January 2022, however, testing was available to all and recommended for even mild symptoms and in contact tracing. Also, individuals with a positive home antigen test result were strongly recommended to confirm this result with another test in the national testing system. Nevertheless, some individuals who had a positive home test result, and asymptomatic individuals who did not test, might have been included in the comparison group, which could reduce the potential associations. Also, a non-random misclassification of infection could have occurred in individuals with hypertension during pregnancy, or in individuals at high risk or with early symptoms of hypertension during pregnancy, because of more intensive testing for SARS-CoV-2 infection in these groups. Increased detection of infection or more negative test results could influence an association in both directions. With the large number of positive test results in the cohort, however, a potential true association would most likely have been seen.

Secondly, from 2020 in Sweden and Norway, proteinuria was no longer compulsory for a diagnosis of pre-eclampsia, in line with the definition of the International Society for the Study of Hypertension

in Pregnancy.³² With the revised definition of pre-eclampsia implemented during the same time as the start of the pandemic, we chose to study hypertension during pregnancy overall and pre-eclampsia specifically, to investigate potential problems of misclassification of pre-eclampsia as gestational hypertension. A rise in incidence of pre-eclampsia after the revised definition could have caused spurious associations between SARS-CoV-2 infection and pre-eclampsia. To limit the possible effect by variation in testing capacity and change in diagnostic criteria over time, we incorporated calendar time in the models by using a stratified Cox model with estimated date of conception as strata, because this method also ensured comparisons were done in individuals at the same gestational ages at a specific calendar time. Also, when pre-eclampsia and gestational hypertension were analysed separately, the effect estimates were similar.

Thirdly, covariates with missing information were managed with a missing indicator, and in a sensitivity analysis by restricting the study period to when information on all covariates was available. Fourthly, although we adjusted for important confounders, such as body mass index, diabetes, and kidney disease, we lacked information on laboratory values and medical chart information describing the symptomatology and severity of disease of the study population. Residual and unmeasured confounding might always be present in observational and registry based studies. Lastly, although our results from two different countries gave similar results, generalisability to other populations with different test strategies and approaches to the pandemic, during a study period of new SARS-CoV-2 variants, or in pregnant individuals with primarily severe covid-19, might be limited.

Conclusion and implications

With nationwide registry data from Sweden and Norway, we did not find any evidence of an association between SARS-CoV-2 infection during pregnancy and subsequent development of hypertension during pregnancy or pre-eclampsia. In this study, we accounted for timing of infection and onset of hypertension during pregnancy. Because infection with the SARS-CoV-2 virus in pregnancy is related to other adverse outcomes of pregnancy, vaccination of pregnant individuals is recommended.

Contributors The study was initiated by AKÖ and OS, and designed by AKÖ, ED, MCM, AS, SEH, and OS. ED and MCM performed the statistical analysis, with support for data analyses by JS. AKÖ wrote the initial draft, with support on interpretation of findings and critical revision by all authors. MCM, SEH, and OS obtained financial support. All authors had full access to the data, and reviewed and approved the final version of the article submitted for publication. AKÖ, ED, MCM, and OS are the guarantors. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. **Transparency:** The lead authors (the manuscript's guarantors) affirm that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any

discrepancies from the study as planned (and, if relevant, registered) have been explained.

Funding The study was supported by the Research Council of Norway (project No 324312), and through its Centres of Excellence funding scheme (project No 262700), and by NordForsk (project No 105545 and No 135876). MCM has received funding from the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement No 947684). The funding sources had no role in the study design; the collection, analysis and interpretation of data; and the writing of the article and the decision to submit it for publication.

Competing interests All authors have completed the ICMJE uniform disclosure form at www.icmje.org/disclosure-of-interest/ and declare: support from the Research Council of Norway and NordForsk for the submitted work; MCM has received funding from the European Research Council under the European Union's Horizon 2020 research and innovation programme; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Ethics approval The study was approved by the Swedish Ethical Review Authority (approval Nos 2020-01499, 2020-02468, 2021-00274) and the Regional Committee for Medical and Health Research Ethics of South/East Norway (No 141135). Each ethics committee provided a waiver of consent for participants because of the use of registry data.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. Data are available by applying to the registry owners at <https://helsedata.no/soknadsveiledning/> and <https://www.medscinet.com/gr/forskare.aspx>. No additional data available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Anne K. Örtqvist <http://orcid.org/0000-0001-9019-9233>

Kari Johansson <http://orcid.org/0000-0002-1497-3079>

Olof Stephansson <http://orcid.org/0000-0003-1528-4563>

REFERENCES

- Gillon TER, Pels A, von Dadelszen P, *et al*. Hypertensive disorders of pregnancy: a systematic review of international clinical practice guidelines. *PLoS One* 2014;9:e113715. [10.1371/journal.pone.0113715](https://doi.org/10.1371/journal.pone.0113715)
- Burton GJ, Redman CW, Roberts JM, *et al*. Pre-Eclampsia: pathophysiology and clinical implications. *BMJ* 2019;366:l2381. [10.1136/bmj.l2381](https://doi.org/10.1136/bmj.l2381)
- Magnus MC, Oakley L, Gjessing HK, *et al*. Pregnancy and risk of COVID-19: a Norwegian registry-linkage study. *BJOG* 2022;129:101–9. [10.1111/1471-0528.16969](https://doi.org/10.1111/1471-0528.16969)
- Gesaka SR, Obimbo MM, Wanyoro A. Coronavirus disease 2019 and the placenta: a literature review. *Placenta* 2022;126:209–23. [10.1016/j.placenta.2022.07.007](https://doi.org/10.1016/j.placenta.2022.07.007)
- Palomo M, Youssef L, Ramos A, *et al*. Differences and similarities in endothelial and angiogenic profiles of preeclampsia and COVID-19 in pregnancy. *Am J Obstet Gynecol* 2022;227:277. [10.1016/j.ajog.2022.03.048](https://doi.org/10.1016/j.ajog.2022.03.048)

- 6 Conde-Agudelo A, Romero R. SARS-cov-2 infection during pregnancy and risk of preeclampsia: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2022;226:68–89. 10.1016/j.ajog.2021.07.009
- 7 Allotey J, Stallings E, Bonet M, *et al.* Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *BMJ* 2020;370:m3320. 10.1136/bmj.m3320
- 8 Papageorgiou AT, Deruelle P, Gunier RB, *et al.* Preeclampsia and COVID-19: results from the INTERCOVID prospective longitudinal study. *Am J Obstet Gynecol* 2021;225:289. 10.1016/j.ajog.2021.05.014
- 9 ERA-EDTA Council, Ortiz A, Cozzolino M, *et al.* Chronic kidney disease is a key risk factor for severe COVID-19: a call to action by the ERA-EDTA. *Nephrol Dial Transplant* 2021;36:87–94. 10.1093/ndt/gfaa314
- 10 Fernandez-Ruiz R, Paredes JL, Niewold TB. COVID-19 in patients with systemic lupus erythematosus: lessons learned from the inflammatory disease. *Transl Res* 2021;232:13–36. 10.1016/j.trsl.2020.12.007
- 11 Rosenbloom JI, Raghuraman N, Carter EB, *et al.* Coronavirus disease 2019 infection and hypertensive disorders of pregnancy. *Am J Obstet Gynecol* 2021;224:623–4. 10.1016/j.ajog.2021.03.001
- 12 Stephansson O, Petersson K, Björk C, *et al.* The Swedish pregnancy register for quality of care improvement and research. *Acta Obstet Gynecol Scand* 2018;97:466–76. 10.1111/aogs.13266
- 13 Norwegian Institute of Public Health. Medical birth registry of Norway. n.d. Available: <https://www.fhi.no/en/hn/health-registries/medical-birth-registry-of-norway/>
- 14 Public Health Agency in Sweden. SmiNet. n.d. Available: <https://www.folkhalsomyndigheten.se/sminet/>
- 15 Norwegian Institute of Public Health. Health registries. n.d. Available: <https://www.fhi.no/en/hn/health-registries/msis/>
- 16 Ludvigsson JF. The first eight months of Sweden's COVID-19 strategy and the key actions and actors that were involved. *Acta Paediatrica* 2020;109:2459–71. 10.1111/apa.15582 Available: <https://onlinelibrary.wiley.com/toc/16512227/109/12>
- 17 Public Health Agency of Sweden. Statistics on SARS-cov-2 variants of special interest. 2022. Available: <https://www.folkhalsomyndigheten.se/smittskydd-beredskap/utbrott/aktuella-utbrott/covid-19/statistik-och-analyser/sars-cov-2-virusvarianter-avsarskild-betydelse/>
- 18 Örtqvist AK, Dahlqvist E, Magnus MC, *et al.* COVID-19 vaccination in pregnant women in Sweden and Norway. *Vaccine* 2022;40:4686–92. 10.1016/j.vaccine.2022.06.083
- 19 Ludvigsson JF, Andersson E, Ekbohm A, *et al.* External review and validation of the Swedish national inpatient register. *BMC Public Health* 2011;11:450. 10.1186/1471-2458-11-450
- 20 Statistics Sweden. Population registers. 2022. Available: <https://www.scb.se/vara-tjanster/bestall-data-och-statistik/bestalla-mikrodata/vilka-mikrodata-finns/>
- 21 Statistics Norway. 2022. Available: <https://www.ssb.no/en>
- 22 Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Stat Med* 2003;22:2693–710. 10.1002/sim.1482
- 23 Gurol-Urganci I, Jardine JE, Carroll F, *et al.* Maternal and perinatal outcomes of pregnant women with SARS-cov-2 infection at the time of birth in England: national cohort study. *Am J Obstet Gynecol* 2021;225:522. 10.1016/j.ajog.2021.05.016
- 24 Jering KS, Claggett BL, Cunningham JW, *et al.* Clinical characteristics and outcomes of hospitalized women giving birth with and without COVID-19. *JAMA Intern Med* 2021;181:714–7. 10.1001/jamainternmed.2020.9241
- 25 HILL AB. The environment and disease: association or causation? *Proc R Soc Med* 1965;58:295–300.
- 26 Shimonovich M, Pearce A, Thomson H, *et al.* Assessing causality in epidemiology: revisiting Bradford Hill to incorporate developments in causal thinking. *Eur J Epidemiol* 2021;36:873–87. 10.1007/s10654-020-00703-7
- 27 Tossetta G, Fantone S, Delli Muti N, *et al.* Preeclampsia and severe acute respiratory syndrome coronavirus 2 infection: a systematic review. *J Hypertens* 2022;40:1629–38. 10.1097/HJH.0000000000003213
- 28 Mendoza M, Garcia-Ruiz I, Maiz N, *et al.* Pre-eclampsia-like syndrome induced by severe COVID-19: a prospective observational study. *BJOG* 2020;127:1374–80. 10.1111/1471-0528.16339
- 29 Birol Ilter P, Prasad S, Mutlu MA, *et al.* Maternal and perinatal outcomes of SARS-cov-2 infection in unvaccinated pregnancies during delta and omicron waves. *Ultrasound Obstet Gynecol* 2022;60:96–102. 10.1002/uog.24916
- 30 Mupanomunda M, Fakhri MG, Miller C, *et al.* Comparison of severe maternal morbidities associated with delivery during periods of circulation of specific SARS-cov-2 variants. *JAMA Netw Open* 2022;5:e2226436. 10.1001/jamanetworkopen.2022.26436
- 31 Seaseley AR, Blanchard CT, Arora N, *et al.* Maternal and perinatal outcomes associated with the omicron variant of severe acute respiratory syndrome coronavirus 2 (SARS-cov-2) infection. *Obstet Gynecol* 2022;140:262–5. 10.1097/AOG.0000000000004849
- 32 Brown MA, Magee LA, Kenny LC, *et al.* Hypertensive disorders of pregnancy: ISSHP classification, diagnosis, and management recommendations for international practice. *Hypertension* 2018;72:24–43. 10.1161/HYPERTENSIONAHA.117.10803
- Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjmed-2022-000465>).

Online Supplement

This supplementary material has been provided by the authors to give readers additional information about their work.

Supplement to: The association between SARS-CoV-2 infection and de novo hypertension during pregnancy – a population-based cohort study

Authors: Anne K Örtqvist, Maria C Magnus, Elisabeth Dahlqvist, Jonas Söderling, Kari Johansson, Anna Sandström, Siri E Håberg, Olof Stephansson

Table of contents	
Data sources and linkages in Sweden	<i>p. 2</i>
Data sources and linkages in Norway	<i>p. 4</i>
Outcome definitions	<i>p. 5</i>
Testing strategies	<i>p. 6</i>
References	<i>p. 7</i>
eTable 1	Participant characteristics according to SARS-CoV-2 infection in 1st, 2nd and 3rd trimester of pregnancy <i>p. 8</i>
eTable 2	Participant characteristics according to SARS-CoV-2 infection during pregnancy for different time periods corresponding to the most dominating virus variant <i>p. 11</i>
eTable 3	Participant characteristics according to hypertension during pregnancy and preeclampsia <i>p. 14</i>
eTable 4	Time between a positive SARS-CoV-2 test and date of diagnosis of hypertension during pregnancy <i>p. 17</i>

Data sources and linkages in Sweden

The Swedish Pregnancy register (SPR)

Data in this study was provided through the Swedish Pregnancy Register (SPR). This quality register was initiated in 2013, and includes 94% of all births in Sweden (18 of 21 regions).(1) Demographical, reproductive and maternal health care data, starting at the first visit to the antenatal care clinic around the ninth gestational week, are transferred from electronic medical records within 24 hours from a reported birth. The register includes information on birth outcomes in addition to maternal background characteristics, health during pregnancy, maternal and neonatal outcomes. From this register we identified the following covariates: gestational age, parity, body mass index at start of pregnancy, smoking during pregnancy, cohabitation with partner, co-existing medical conditions and the outcome of hypertension during pregnancy.

Registers at the National Board of Health and Welfare

Covariates related to prior health conditions according to the International Classification of Diseases tenth revision (ICD-10) diagnose codes and/or Anatomical Therapeutic Chemical (ATC) codes for pre-existing hypertension and debut of hypertension up till gestational week 20, pre-existing diabetes, chronic kidney disease, and systemic lupus erythematosus, as well as number of hospital visits and stays were also collected from the National Patient Register(2) and the Prescribed Drug Register(3) (March 1, 2020-October 10, 2021) with a lookback of 3 years from March 2020.

Registers at the Public Health Agency of Sweden

From the start of the pandemic in March 2020, the SPR has been linked to the Swedish Register for Communicable Diseases (SmiNet) and the national vaccination register to retrieve regularly updated data on PCR-verified SARS-CoV-2 tests and vaccination coverage among all birthing women in the SPR.

SmiNet is the national register for communicable diseases. On February 1, 2020, SARS-CoV-2 was included in the Swedish Communicable Diseases Act, making it mandatory to report all laboratory-confirmed Polymerase Chain Reaction (PCR) cases within 24 hours to the register. From this register we used information on positive tests as well as the date of a positive PCR test. Information on key elements regarding the Swedish pandemic strategy has been discussed in detail papers by Ludvigsson JF.(4, 5)

As of January 1, 2013, healthcare providers must report all vaccinations administered within the Swedish vaccination programs to the Swedish vaccination register, held by the Public Health Agency of Sweden. The register includes type and date of all COVID-19 vaccinations. Up to end of May, 2021 in Sweden, vaccination was only recommended to pregnant women with a high risk of severe COVID-19. Thereafter, a general recommendation for pregnant women to get vaccinated was issued, although women were recommended to wait with vaccinations to after 12 weeks of gestation. As availability of vaccines was restricted initially, vaccination was still being prioritized based on age (oldest first). From August, 2021, however, the vaccine was available to all above 18 years of age. More information on COVID-19 vaccinations among pregnant and birthing women in Sweden and Norway can be found in recent publications by Örtqvist AK et al.(6) and Magnus MC et al.(7)

Registers at Statistics Sweden

Education, income (average individual income from work during 2017-2019 in tertiles) and country of birth were collected from the Total Population Register, Education Register and Income and tax register at Statistics Sweden (March 1, 2020-October 10, 2021).(8) Information on educational level and country of birth, for the period October 11, 2021-May 24, 2022, were also collected from the Swedish Pregnancy Register as this time period was not covered by the information from Statistics Sweden.

Data sources and linkages in Norway

The Emergency Preparedness Register for Covid-19

Data in this study were provided through the Emergency preparedness register for Covid-19 (Beredt C19) administered by the Norwegian Institute of Public Health, according to the Health Preparedness Act §2-4.(9) This registry was established in 2020 to provide authorities with up to date information on prevalence, causal relationships, and consequences of the Covid19 epidemic in Norway. Beredt C19 includes information already collected in the healthcare service, national health registries and administrative registers with information about the Norwegian population. The data subjects' right is safeguarded as they can contact the data controller for all different sources included in Beredt C19 in the usual way. Through Beredt C19 we used data from the following sources:

The Medical Birth Registry of Norway (MBRN)

The Norwegian Medical Birth Registry includes information on all pregnancies ending in gestational week 12 or later.(10) The registry includes information on birth outcomes in addition to maternal background characteristics, health during pregnancy, pregnancy outcomes and neonatal health.

National Patient Register

The Norwegian patient registry includes individual level information on all contacts with specialist health-care services. Information registered includes admission and discharge dates, and diagnostic codes during the hospital stay. These discharge codes are coded according to the International Classification of Diseases version 10. From this registry information on pre-existing hypertension, co-existing medical conditions and the outcomes of hypertension during pregnancy were identified.

Norwegian Surveillance System for Communicable Diseases (MSIS)

There is mandatory reporting of selected infectious diseases to this surveillance register.(11) Reporting of all Covid-19 tests is mandatory, and this register contains date of testing and test results.

The Norwegian Immunisation Register (SYSVAK)

SYSVAK is a register of vaccines in the Norwegian vaccination program, with mandatory registrations of all Covid-19 vaccinations (dates and type). Up to end of August, 2021 in Norway, vaccination was only recommended to pregnant women with a high risk of severe COVID-19. Thereafter, a general recommendation for pregnant women to get vaccinated was issued, although women were recommended to wait with vaccinations to after 12 weeks of gestation. Since January, 2022, women in Norway have been recommended vaccination regardless of trimester.(6, 7)

Statistics Norway (SSB)

Administrative data is mandatorily reported to Statistics Norway.(12) We used information from this database on household income in 2018, type of education and years of education completed by 2019.

Definition of hypertension during pregnancy (HDP) and date of diagnosis (used as proxy for disease onset) based from the national registers

HDP (excluding pre-existing hypertension or hypertension during pregnancy up until gestational week 20*) was defined by International Classification of Disease (ICD)-10 codes: O13.9, O14 or O15 identified from the national patient registries respectively (inpatient or outpatient care) in Sweden and Norway, from gestational week 20+0 up until 1 week after delivery (to identify possible postpartum preeclampsia) or the SPR and MBRN (diagnosis registered at delivery).

The following criteria was applied:

- (1) one inpatient care admission, or
- (2) two outpatient visits, or
- (3) one outpatient visit followed by a one inpatient care admission, or
- (4) one outpatient visit followed by a diagnosis in the SPR/MBRN, or
- (5) a diagnosis only in the SPR/MBRN.

The date of preeclampsia diagnosis was defined as the first recording in inpatient care, outpatient care or the SPR/MBRN. Since factors (hypertension) determining diagnosis of gestational hypertension are also first signs of preeclampsia we did the following reclassification: If date of gestational hypertension occurred after diagnosis of preeclampsia this was regarded as misclassification of preeclampsia for gestational hypertension. If gestational hypertension occurred prior to diagnosis of preeclampsia we regarded this as a first indication of preeclampsia and used the date of diagnosis of gestational hypertension as date of diagnosis of preeclampsia. For women with no registration of hypertension disorders of pregnancy in the Norwegian patient registry, and the only registration was observed in the MBRN, the onset of hypertensive disorders of pregnancy was defined to be the gestational day of delivery.

***Definition of pre-existing hypertension**

Pre-existing hypertension and hypertension during pregnancy up until gestational week 20, was defined based on ICD-10 codes O10 and/or I10-I15 in the national patient registries, SPR/MBRN, and/or from Anatomical Therapeutic Chemical (ATC) codes (C02, C03, C07, C08, C09) in the Swedish Prescribed Drug Register.

Testing strategies

In Sweden, SARS-CoV-2 was included in the Swedish Communicable Disease Act on 1st February, 2020, making all laboratory-confirmed polymerase chain reaction (PCR) cases of SARS-CoV-2 mandatory to report within 24 hours to SmiNet at the Public Health Agency of Sweden. Negative tests results are unfortunately not available on a national level. A non-universal population testing strategy was implemented including outpatient testing and contact tracing, starting in June 2020 and ongoing to January 2022. This type of testing mostly included symptomatic individuals, but it could also include individuals tested before and after travels, or after contact with other test-positive individuals and individuals subject to workplace testing (e.g. healthcare workers). In 23 of the 39 delivery hospitals covered by the Swedish Pregnancy Register, universal testing of all women admitted for labor or pregnancy in-patient care, independent of their current and previous medical history and COVID-19 symptoms or not, were performed.

In Norway, mandatory registration of all PCR tests for SARS-CoV-2 was implemented in the Norwegian Surveillance System for Communicable Diseases on 31st January, 2020. As Norway has not had universal testing of pregnant or delivering women, a non-universal testing strategy was operating throughout the study period. Testing was predominantly conducted on the basis of symptoms to confirm or exclude SARS-CoV-2 infection, with some additional testing conducted for particular reasons such as contact with infected persons, or mandatory testing due to travel or work. Information was available on all conducted tests, regardless of positive or negative result.

From mid-January 2022, the testing has decreased due to new recommendations. Individuals are no longer required to get tested, but to stay at home if one has symptoms. However, healthcare workers, and pregnant women with symptoms of COVID-19 in contact with health care, are still recommended to get tested, and several delivery hospitals still test pregnant women upon admission to the delivery ward independent of symptoms.

With regards to the varying testing strategies in the two countries over time, we performed a stratified cox model which compares individuals with the same estimated date of conception with each other.

References

1. Stephansson O, Petersson K, Bjork C, Conner P, Wikstrom AK. The Swedish Pregnancy Register - for quality of care improvement and research. *Acta obstetrica et gynecologica Scandinavica*. 2018;97(4):466-76.
2. Ludvigsson JF, Andersson E, Ekblom A, Feychting M, Kim JL, Reuterwall C, et al. External review and validation of the Swedish national inpatient register. *BMC Public Health*. 2011;11:450.
3. Wallerstedt SM, Wettermark B, Hoffmann M. The First Decade with the Swedish Prescribed Drug Register - A Systematic Review of the Output in the Scientific Literature. *Basic & clinical pharmacology & toxicology*. 2016;119(5):464-9.
4. Ludvigsson JF. The first eight months of Sweden's COVID-19 strategy and the key actions and actors that were involved. *Acta paediatrica (Oslo, Norway : 1992)*. 2020;109(12):2459-71.
5. Ludvigsson JF. How Sweden approached the COVID-19 pandemic: summary and commentary on the national commission inquiry. *Acta paediatrica (Oslo, Norway : 1992)*. 2022.
6. Ortqvist AK, Dahlqvist E, Magnus MC, Ljung R, Jonsson J, Aronsson B, et al. COVID-19 vaccination in pregnant women in Sweden and Norway. *Vaccine*. 2022;40(33):4686-92.
7. Magnus MC, Ortqvist AK, Dahlqvist E, Ljung R, Skar F, Oakley L, et al. Association of SARS-CoV-2 Vaccination During Pregnancy With Pregnancy Outcomes. *Jama*. 2022;327(15):1469-77.
8. Statistics Sweden. Population registers [cited 2022 Oct 4]. Available from: <https://www.scb.se/vara-tjanster/bestall-data-och-statistik/bestalla-mikrodata/vilka-mikrodata-finns/>.
9. Norwegian Institute of Public Health. Emergency preparedness register for COVID-19 [cited 2022 Oct 4]. Available from: <https://www.fhi.no/en/id/infectious-diseases/coronavirus/emergency-preparedness-register-for-covid-19/>.
10. Norwegian Institute of Public Health. Medical Birth Registry of Norway [cited 2022 18 September]. Available from: <https://www.fhi.no/en/hn/health-registries/medical-birth-registry-of-norway/>.
11. Norwegian Institute of Public Health. Norwegian Surveillance System for Communicable Diseases [cited 2022 Oct 4]. Available from: <https://www.fhi.no/en/hn/health-registries/msis/>.
12. Statistics Norway [cited 2022 Oct 4]. Available from: <https://www.ssb.no/en>.

eTable 1. Participant characteristics according to SARS-CoV-2 infection in 1st, 2nd and 3rd trimester of pregnancy

	Sweden				Norway			
	Not SARS-CoV-2	SARS-CoV-2			Not SARS-CoV-2	SARS-CoV-2		
		1 st trimester	2 nd trimester	3 rd trimester		1 st trimester	2 nd trimester	3 rd trimester
N	182 810 (100.0)	3260 (100.0)	6480 (100.0)	9219 (100.0)	105 090 (100.0)	428 (100.0)	1257 (100.0)	3911 (100.0)
CHARACTERISTICS								
SARS-CoV-2 vaccination - n (%)								
Vaccinated prior pregnancy	5787 (3.2)	55 (1.7)	787 (12.1)	997 (10.8)	643 (0.6)	18 (4.2)	131 (10.4)	257 (6.6)
Vaccinated dose 1 during pregnancy	27 533 (15.1)	876 (26.9)	844 (13.0)	2563 (27.8)	15 586 (14.8)	86 (20.1)	396 (31.2)	265 (57.9)
Not vaccinated	149 490 (81.8)	2329 (71.4)	4849 (74.8)	5659 (61.4)	88 861 (84.6)	324 (75.7)	730 (58.1)	1389 (35.5)
Maternal age (years) - n (%)								
<20	2046 (1.1)	10 (0.3)	40 (0.6)	99 (1.1)	902 (0.9)	7 (1.6)	9 (0.7)	20 (0.5)
20-24	18 336 (10.0)	324 (9.9)	635 (9.8)	894 (9.7)	10 575 (10.1)	47 (11.0)	108 (8.6)	323 (8.3)
25-29	60 887 (33.3)	1136 (34.8)	2201 (34.0)	3061 (33.2)	35 796 (34.1)	122 (28.5)	396 (31.5)	1318 (33.7)
30-34	66 754 (36.5)	1188 (36.4)	2418 (37.3)	3372 (36.6)	39 055 (37.2)	159 (37.2)	498 (39.6)	1540 (39.4)
35-39	28 993 (15.9)	481 (14.8)	992 (15.3)	1498 (16.2)	15 992 (15.2)	82 (19.2)	201 (16.0)	616 (15.8)
>40	5794 (3.2)	121 (3.7)	194 (3.0)	295 (3.2)	2770 (2.6)	11 (2.6)	45 (3.6)	94 (2.4)
Body Mass Index - n (%)								
<18.5	4083 (2.2)	47 (1.4)	112 (1.7)	171 (1.9)	3253 (3.1)	10 (2.3)	31 (2.5)	117 (3.0)
18.5-<25	94 922 (51.9)	1605 (49.2)	3241 (50.0)	4501 (48.8)	57 759 (55.0)	222 (51.9)	712 (56.6)	2132 (54.5)
25-<30	48 708 (26.6)	916 (28.1)	1810 (27.9)	2648 (28.7)	23 068 (22.0)	105 (24.5)	276 (22.0)	918 (23.5)
>30	28 039 (15.3)	586 (18.0)	1092 (16.9)	1586 (17.2)	13 765 (13.1)	64 (15.0)	159 (12.7)	514 (13.1)
Missing	058 (3.9)	106 (3.3)	225 (3.5)	313 (3.4)	7245 (6.9)	27 (6.3)	79 (6.3)	230 (5.9)
Parity - n (%)								
Parity 0	79 398 (43.4)	1352 (41.5)	2617 (40.4)	3526 (38.2)	46 277 (44.0)	172 (40.2)	460 (36.6)	1460 (37.3)
Parity 1	67 885 (37.1)	1180 (36.2)	2458 (37.9)	3505 (38.0)	38 660 (36.8)	135 (31.2)	435 (34.6)	1491 (38.1)
Parity 2 or more	35 527 (19.4)	728 (22.3)	1405 (21.7)	2188 (23.7)	20 153 (19.2)	121 (28.3)	362 (28.8)	960 (24.6)
Educational level (years) - n (%)								
<9	16 812 (9.2)	224 (6.9)	477 (7.4)	763 (8.3)	13 722 (10.1)	85 (19.9)	251 (20.0)	569 (14.6)
10-12	66 543 (36.4)	1289 (39.5)	2414 (37.3)	3144 (34.1)	20 138 (19.2)	84 (9.6)	245 (19.5)	766 (19.6)
>12	89 896 (49.2)	1551 (47.6)	3177 (49.0)	4338 (47.1)	61 255 (58.3)	179 (41.8)	573(45.6)	2126 (54.4)

eTable 1. Cont.	Sweden				Norway			
	Not SARS-CoV-2	SARS-CoV-2			Not SARS-CoV-2	SARS-CoV-2		
		1 st trimester	2 nd trimester	3 rd trimester		1 st trimester	2 nd trimester	3 rd trimester
<i>Education level, cont.</i>								
<i>Missing</i>	9559 (5.2)	196 (6.0)	412 (6.4)	974 (10.6)	9975 (9.5)	80 (18.7)	188 (15.0)	450 (11.5)
Income in tertiles- n (%)								
Tertile 1	46 717 (25.6)	606 (18.6)	1329 (20.5)	1429 (15.5)	45 898 (43.7)	233 (54.4)	646 (51.4)	1879 (48.0)
Tertile 2	46 548 (25.5)	686 (21.0)	1587 (24.5)	1258 (13.6)	30 324 (28.9)	102 (23.8)	289 (23.0)	1022 (26.1)
Tertile 3	46 859 (25.6)	645 (19.8)	1388 (21.4)	1185 (12.9)	23 497 (22.4)	64 (15.0)	227 (18.1)	783 (20.0)
<i>Missing</i>	40 387 (22.1)	1302 (39.9)	2135 (32.9)	5264 (57.1)	5371 (5.1)	29 (6.8)	95 (7.6)	227 (5.8)
Birth region - n (%)								
Scandinavia	126 378 (69.1)	2304 (70.7)	4459 (68.8)	5992 (65.0)	78 400 (74.6)	228 (53.3)	682 (54.3)	2703 (69.1)
Other European countries	14 574 (8.0)	253 (7.8)	539 (8.3)	709 (7.7)	11 728 (11.2)	69 (16.1)	214 (17.0)	484 (12.4)
Middle East/ Africa	28 826 (15.8)	504 (15.5)	1025 (15.8)	1677 (18.2)	6987 (6.7)	91 (21.3)	244 (19.4)	429 (11.0)
Other	8518 (4.7)	115 (3.5)	219 (3.4)	325 (3.5)	7960 (7.6)	40 (9.4)	117 (9.3)	294 (7.5)
<i>Missing</i>	4514 (2.5)	84 (2.6)	238 (3.7)	516 (5.6)	15 (0.01)	0	0	1 (0.03)
Smoking status - n (%)								
Non-smoker (reference)	170 485 (93.3)	3064 (94.0)	6072 (93.7)	8588 (93.2)	89 050 (84.7)	345 (80.6)	1027 (81.7)	3306 (84.5)
Smoker	6101 (3.3)	80 (2.5)	177 (2.7)	276 (3.0)	5269 (5.0)	23 (5.4)	71 (5.7)	169 (4.3)
<i>Missing</i>	6224 (3.4)	116 (3.6)	231 (3.6)	355 (3.9)	10 771 (10.3)	60 (14.0)	159 (12.7)	436 (11.2)
Living with partner - n (%)								
Yes	165 110 (90.3)	3020 (92.6)	5964 (92.0)	8351 (90.6)	99 197 (94.4)	385 (90.0)	1163 (92.5)	3700 (94.6)
No	13 727 (7.5)	179 (5.5)	402 (6.2)	690 (7.5)	4153 (4.0)	32 (7.5)	74 (5.9)	152 (3.9)
<i>Missing</i>	3973 (2.2)	61 (1.9)	114 (1.8)	178 (1.9)	1740 (1.7)	11 (2.6)	20 (1.6)	59 (1.5)
Co-existing medical conditions - n (%)								
Gestational diabetes	8184 (4.5)	159 (4.9)	316 (4.9)	370 (4.0)	6802 (6.5)	36 (8.4)	102 (8.1)	241 (6.2)
Pre-existing diabetes mellitus	1994 (1.1)	48 (1.5)	84 (1.3)	98 (1.1)	790 (0.8)	3 (0.7)	5 (0.4)	23 (0.6)
Chronic kidney disease	645 (0.4)	12 (0.4)	26 (0.4)	28 (0.3)	575 (0.6)	2 (0.5)	8 (0.6)	23 (0.6)

eTable 1. Cont.	Sweden				Norway			
	Not SARS-CoV-2	SARS-CoV-2			Not SARS-CoV-2	SARS-CoV-2		
		1 st trimester	2 nd trimester	3 rd trimester		1 st trimester	2 nd trimester	3 rd trimester
<i>Co-existing medical conditions, cont.</i>								
Systemic Lupus Erythematosus	259 (0.1)	3 (0.1)	13 (0.2)	14 (0.2)	88 (0.1)	0	0	2 (0.1)
History of hypertension during pregnancy - n (%)								
Yes	1413 (0.8)	20 (0.6)	59 (0.9)	33 (0.4)	3517 (3.4)	10 (2.3)	47 (3.7)	142 (3.6)
Missing	40 387 (22.1)	1302 (39.9)	2135 (32.9)	5264 (57.1)	0	0	0	0
Visits to outpatient care 1 year prior								
0 visits	87 900 (48.1)	1181 (36.2)	2629 (40.6)	2421 (26.3)	n/a	n/a	n/a	n/a
1-9 visits	51 839 (28.4)	745 (22.9)	1644 (25.4)	1469 (15.9)	n/a	n/a	n/a	n/a
10-29 visits	2637 (1.4)	30 (0.9)	71 (1.1)	63 (0.7)	n/a	n/a	n/a	n/a
30 + visits	47 (0.0)	2 (0.1)	1 (0.0)	2 (0.0)	n/a	n/a	n/a	n/a
Missing	40 387 (22.1)	1302 (39.9)	2135 (32.9)	5264 (57.1)	105 090 (100.0)	428 (100.0)	1257 (100.0)	3911 (100.0)
Gestational age at birth - n (%)								
Week 22+0 -36+6	7686 (4.2)	108 (3.3)	289 (4.5)	402 (4.4)	4726 (4.5)	17 (4.0)	56 (4.5)	122 (3.1)
Week 37+0-39+6	83 388 (45.6)	1480 (45.4)	2953 (45.6)	4353 (47.2)	43 972 (41.8)	179 (41.8)	572 (45.5)	1664 (42.6)
Week >= 40+0	91 736 (50.2)	1672 (51.3)	3238 (50.0)	4464 (48.4)	56 392 (53.7)	232 (54.2)	629 (50.0)	2125 (54.3)
Outcomes - n (%)								
Hypertension during pregnancy	10 591 (5.8)	175 (5.4)	365 (5.6)	489 (5.3)	6122 (5.8)	18 (4.2)	69 (5.5)	222 (5.7)
Preeclampsia	5390 (2.9)	90 (2.8)	174 (2.7)	237 (2.6)	3822 (3.6)	13 (3.0)	43 (3.4)	130 (3.3)

eTable 2. Participant characteristics according to SARS-CoV-2 during pregnancy for different time periods corresponding to the most dominating virus variant

	Sweden				Norway			
	Index	Alpha	Delta	Omicron	Index	Alpha	Delta	Omicron
N	6655 (100.0)	4855 (100.0)	1744 (100.0)	5706 (100.0)	574 (100.0)	590 (100.0)	1221 (100.0)	3211 (100.0)
CHARACTERISTICS								
SARS-CoV-2 vaccination - n (%)								
Vaccinated prior pregnancy	0	23 (0.5)	248 (14.2)	1568 (27.5)	0	4 (0.7)	116 (9.5)	286 (8.9)
Vaccinated dose 1 during pregnancy	326 (4.9)	908 (18.7)	569 (32.6)	2481 (43.5)	7 (1.2)	62 (10.5)	420 (34.4)	2258 (70.3)
Not vaccinated	6329 (95.1)	3924 (80.8)	927 (53.2)	1657 (29.0)	567 (98.8)	524 (88.8)	685 (56.1)	667 (20.8)
Maternal age (years) - n (%)								
<20	42 (0.6)	28 (0.6)	25 (1.4)	54 (0.9)	2 (0.4)	7 (1.2)	15 (1.2)	12 (0.4)
20-24	698 (10.5)	480 (9.9)	171 (9.8)	503 (8.8)	59 (10.3)	66 (11.2)	104 (8.5)	249 (7.8)
25-29	2312 (34.7)	1690 (34.8)	554 (31.8)	1840 (32.2)	194 (33.8)	175 (29.7)	378 (31.0)	1089 (33.9)
30-34	2400 (36.1)	1790 (36.9)	615 (35.3)	2175 (38.1)	223 (38.9)	233 (39.5)	461 (37.8)	1280 (39.9)
35-39	1002 (15.1)	704 (14.5)	317 (18.2)	949 (16.6)	79 (13.8)	88 (14.9)	225 (18.4)	507 (15.8)
>40	201 (3.0)	163 (3.4)	62 (3.6)	185 (3.2)	17 (3.0)	21 (3.6)	38 (3.1)	74 (2.3)
Body Mass Index - n (%)								
<18.5	101 (1.5)	82 (1.7)	39 (2.2)	108 (1.9)	14 (2.4)	14 (2.4)	39 (3.2)	91 (2.8)
18.5-<25	3282 (49.3)	2423 (49.9)	814 (46.7)	2826 (49.5)	294 (51.2)	295 (50.0)	679 (55.6)	1798 (56.0)
25-<30	1864 (28.0)	1382 (28.5)	528 (30.3)	1601 (28.1)	148 (25.8)	135 (22.9)	273 (22.4)	743 (23.1)
>30	1182 (17.8)	810 (16.7)	299 (17.1)	975 (17.1)	75 (13.1)	96 (16.3)	159 (13.0)	407 (12.7)
Missing	226 (3.4)	158 (3.3)	64 (3.7)	196 (3.4)	43 (7.5)	50 (8.5)	71 (5.8)	172 (5.4)
Parity - n (%)								
Parity 0	2828 (42.5)	1896 (39.1)	686 (39.3)	2086 (36.6)	213 (37.1)	207 (35.1)	484 (39.6)	1188 (37.0)
Parity 1	2375 (35.7)	1885 (38.8)	591 (33.9)	2291 (40.2)	196 (34.2)	221 (37.5)	399 (32.7)	1245 (38.8)
Parity 2 or more	1452 (21.8)	1074 (22.1)	467 (26.8)	1329 (23.3)	165 (28.8)	162 (27.5)	338 (27.7)	778 (24.2)
Educational level (years) - n (%)								
<9	645 (9.7)	397 (8.2)	139 (8.0)	284 (5.0)	115 (20.0)	122 (20.7)	262 (21.5)	406 (12.6)

eTable 2. Cont.	Sweden				Norway			
	Index	Alpha	Delta	Omicron	Index	Alpha	Delta	Omicron
<i>Education level, cont.</i>								
10-12	2726 (41.0)	1957 (40.3)	535 (30.7)	1627 (28.5)	107 (18.6)	122 (20.7)	216 (17.7)	650 (20.2)
>12	3209 (48.2)	2286 (47.1)	774 (44.4)	2800 (49.1)	252 (43.9)	237 (40.2)	554 (45.4)	1835 (57.2)
<i>Missing</i>	75 (1.1)	215 (4.4)	296 (17.0)	995 (17.4)	100 (17.4)	109 (18.5)	189 (15.5)	320 (10.0)
Income in tertiles - n (%)								
Tertile 1	2161 (32.5)	1154 (23.8)	48 (2.8)	0	225 (39.2)	9 (40.5)	496 (40.6)	1077 (33.5)
Tertile 2	2270 (34.1)	1234 (25.4)	27 (1.5)	0	175 (30.5)	166 (28.1)	351 (38.8)	1010 (31.5)
Tertile 3	2103 (31.6)	1095 (22.6)	22 (1.3)	0	138 (24.0)	143 (24.2)	283 (23.2)	942 (29.3)
<i>Missing</i>	54 (0.8)	1302 (26.8)	1638 (93.9)	5706 (100)	36 (6.3)	42 (7.1)	91 (7.5)	182 (5.7)
Birth region - n (%)								
Scandinavia	4460 (67.0)	3462 (71.3)	993 (56.9)	3840 (67.3)	294 (51.2)	288 (48.8)	666 (54.6)	2365 (73.7)
Other European countries	585 (8.8)	372 (7.7)	169 (9.7)	374 (6.6)	80 (13.9)	107 (18.1)	255 (18.4)	355 (11.1)
Middle East/ Africa	1356 (20.4)	755 (15.6)	343 (19.7)	751 (13.2)	143 (24.9)	139 (23.6)	213 (17.4)	269 (8.4)
Other	248 (3.7)	181 (3.7)	61 (3.5)	171 (3.0)	56 (9.8)	56 (9.5)	117 (9.6)	222 (6.9)
<i>Missing</i>	6 (0.1)	85 (1.8)	178 (10.2)	570 (10.0)	1 (0.2)	0	0	0
Smoking status - n (%)								
Non-smoker (reference)	6252 (93.9)	4565 (94.0)	1601 (91.8)	5307 (93.0)	473 (82.4)	480 (81.4)	999 (81.8)	2726 (84.9)
Smoker	172 (2.6)	129 (2.7)	66 (3.8)	166 (2.9)	36 (6.3)	32 (5.4)	60 (4.9)	135 (10.9)
<i>Missing</i>	231 (3.5)	161 (3.3)	77 (4.4)	233 (4.1)	65 (11.3)	78 (13.2)	162 (13.3)	350 (10.9)
Living with partner - n (%)								
Yes	6122 (92.0)	4456 (91.8)	1576 (90.4)	5183 (90.8)	534 (93.0)	528 (89.5)	1131 (92.6)	3055 (95.1)
No	427 (6.4)	298 (6.1)	137 (7.9)	408 (7.2)	29 (5.1)	44 (7.5)	72 (5.9)	113 (3.5)
<i>Missing</i>	106 (1.6)	101 (2.1)	31 (1.8)	115 (2.0)	11 (1.9)	18 (3.1)	18 (1.5)	43 (1.3)
Co-existing medical conditions - n (%)								
Gestational diabetes	441 (6.6)	235 (4.8)	55 (3.2)	115 (2.0)	46 (8.0)	52 (8.9)	102 (8.4)	179 (5.6)
Pre-existing diabetes mellitus	84 (1.3)	64 (1.3)	20 (1.1)	61 (1.1)	6 (1.1)	2 (0.3)	6 (0.5)	17 (0.5)
Chronic kidney disease	28 (0.4)	20 (0.4)	3 (0.2)	15 (0.3)	1 (0.2)	1 (0.2)	11 (0.9)	20 (0.6)

eTable 2. Cont.	Sweden				Norway			
<i>Co-existing medical conditions, cont.</i>	Index	Alpha	Delta	Omicron	Index	Alpha	Delta	Omicron
Systemic Lupus Erythematosus	9 (0.1)	10 (0.2)	0	11 (0.2)	1 (0.2)	0	0	1 (0.03)
History of hypertension during pregnancy- n (%)								
Yes	69 (1.0)	42 (0.9)	1 (0.1)	0	15 (2.6)	23 (3.9)	32 (2.6)	129 (4.0)
<i>Missing</i>	54 (0.8)	1302 (26.8)	1638 (93.9)	5706 (100)	0	0	0	0
Visits to outpatient care 1 year prior - n (%)								
0 visits	3972 (59.7)	2192 (45.1)	68 (3.9)	n/a	n/a	n/a	n/a	n/a
1-9 visits	2515 (37.8)	1307 (26.9)	37 (2.1)	n/a	n/a	n/a	n/a	n/a
10-29 visits	112 (1.7)	51 (1.1)	1 (0.1)	n/a	n/a	n/a	n/a	n/a
30 + visits	2 (0.0)	3 (0.1)	(0.0)	n/a	n/a	n/a	n/a	n/a
<i>Missing</i>	54 (0.8)	1302 (26.8)	1638 (93.9)	5706 (100)	574 (100.0)	590 (100.0)	1221 (100.0)	3211 (100.0)
Gestational age at birth - n (%)								
Week 22+0 -36+6	326 (4.9)	207 (4.3)	75 (4.3)	188 (3.3)	23 (4.0)	18 (3.1)	46 (3.8)	108 (3.4)
Week 37+0-39+6	3068 (46.1)	2178 (44.9)	811 (46.5)	2 732 (47.9)	234 (40.8)	274 (46.4)	548 (44.9)	1359 (42.3)
Week >= 40+0	3261 (49.0)	2470 (50.9)	858 (49.2)	2 786 (48.8)	317 (55.2)	298 (50.5)	627 (51.4)	1744 (54.3)
Outcomes - n (%)								
Hypertension during pregnancy	429 (6.4)	245 (5.0)	73 (4.2)	282 (4.9)	22 (3.8)	25 (4.2)	66 (5.4)	196 (6.1)
Preeclampsia	231 (3.5)	121 (2.5)	33 (1.9)	116 (2.0)	13 (2.3)	13 (2.2)	42 (3.4)	118 (3.7)

^aThe background characteristics for those without SARS-CoV-2 (Sweden: n=182 810; Norway: n=105 090) can be found in eTable 1. Overall, those with no positive test and those with a positive test in each time period, corresponds to the total population of 201 770 in Sweden and 110 686 in Norway.

eTable 3. Participant characteristics according to hypertension during pregnancy and preeclampsia

	Sweden			Norway		
	No hypertension during pregnancy	Hypertension during pregnancy	Preeclampsia	No hypertension during pregnancy	Hypertension during pregnancy	Preeclampsia
N	190 150 (100.0)	11 620 (100.0)	5891 (100.0)	104 255 (100.0)	6431 (100.0)	4008 (100.0)
CHARACTERISTICS						
SARS-CoV-2 during pregnancy - n (%)						
Yes	17 931 (9.4)	1029 (8.9)	501 (8.5)	5287 (5.1)	309 (4.8)	186 (4.6)
No	172 219 (90.6)	10 591 (91.1)	5390 (91.5)	98 968 (94.9)	6122 (95.2)	3822 (95.4)
COVID-19 vaccination - n (%)						
Vaccinated prior pregnancy	7124 (3.7)	502 (4.3)	231 (3.9)	975 (0.9)	74 (1.2)	50 (1.3)
Vaccinated dose 1 during pregnancy	29 922 (15.7)	1895 (16.3)	890 (15.1)	17 064 (16.4)	1269 (19.7)	780 (19.7)
Not vaccinated	153 104 (80.5)	9223 (79.4)	4770 (81.0)	82 216 (82.7)	5088 (79.1)	3169 (79.1)
Maternal age (years) - n (%)						
<20	2071 (1.1)	124 (1.1)	87 (1.5)	851 (0.8)	87 (1.4)	75 (1.9)
20-24	19 016 (10.0)	1172 (10.1)	671 (11.4)	10 242 (9.8)	811 (12.6)	567 (14.2)
25-29	63 288 (33.3)	3995 (34.4)	2069 (35.1)	35 433 (34.0)	2199 (34.2)	1382 (34.5)
30-34	69 750 (36.7)	3984 (34.3)	1968 (33.4)	39 105 (37.5)	2147 (33.4)	1287 (32.1)
35-39	30 122 (15.8)	1843 (15.9)	876 (14.9)	15 919 (15.3)	972 (15.1)	573 (14.3)
>40	5903 (3.1)	502 (4.3)	220 (3.7)	2705 (2.6)	215 (3.3)	124 (3.1)
Body Mass Index - n (%)						
<18.5	4278 (2.2)	135 (1.2)	87 (1.5)	3296 (3.2)	115 (1.8)	91 (2.3)
18.5-<25	99 903 (52.5)	4364 (37.6)	2298 (39.0)	58 146 (55.8)	2679 (41.7)	1710 (42.7)
25-<30	50 503 (26.6)	3580 (30.8)	1776 (30.1)	22650 (21.7)	1717 (26.7)	1046 (26.1)
>30	28 142 (14.8)	3163 (27.2)	1515 (25.7)	12938 (12.4)	1564 (24.3)	941 (23.5)
Missing	7324 (3.9)	378 (3.3)	215 (3.6)	7225 (6.9)	356 (5.5)	220 (5.5)
Parity - n (%)						
Parity 0	79 230 (41.7)	7664 (66.0)	4136 (70.2)	44 315 (42.5)	4054 (63.0)	2643 (65.9)

eTable 3. Cont.	Sweden			Norway		
	No hypertension during pregnancy	Hypertension during pregnancy	Preeclampsia	No hypertension during pregnancy	Hypertension during pregnancy	Preeclampsia
<i>Parity, cont.</i>						
Parity 1	72 397 (38.1)	2630 (22.6)	1158 (19.7)	39 124 (37.5)	1597 (24.8)	891 (22.2)
Parity 2 or more	38 523 (20.3)	1326 (11.4)	597 (10.1)	20 816 (20.0)	780 (12.1)	474 (11.8)
Educational level (years) - n (%)						
<9	17 483 (9.2)	794 (6.8)	484 (8.2)	13 681 (13.1)	946 (14.7)	654 (6.3)
10-12	68 913 (36.2)	4475 (38.5)	2270 (38.5)	19 817 (19.0)	1416 (22.0)	926 (23.1)
>12	93 123 (49.0)	5842 (50.3)	2866 (48.7)	60 496 (58.0)	3637 (56.6)	2150 (53.6)
<i>Missing</i>	10 631 (5.6)	509 (4.4)	271 (4.6)	10 261 (9.8)	432 (6.7)	278 (6.9)
Income in tertiles - n (%)						
Tertile 1	47 613 (25.0)	2467 (21.2)	1400 (23.8)	32 750 (31.4)	2238 (34.8)	1460 (36.4)
Tertile 2	47 139 (24.8)	2940 (25.3)	1462 (24.8)	32 982 (31.6)	2006 (31.2)	1234 (30.8)
Tertile 3	46 583 (24.5)	3496 (30.1)	1689 (28.7)	33 047 (31.7)	1941 (30.2)	1150 (28.7)
<i>Missing</i>	46 451 (24.4)	2636 (22.7)	1287 (21.8)	5476 (5.3)	246 (3.8)	164 (4.1)
Birth region - n (%)						
Scandinavia	129 898 (68.3)	9235 (79.5)	4493 (76.3)	76 763 (73.6)	5250 (81.6)	3232 (80.6)
Other European countries	15 330 (8.1)	744 (6.4)	369 (6.3)	11 950 (11.5)	545 (8.5)	327 (8.2)
Middle East/ Africa	31 059 (16.3)	972 (8.4)	641 (10.9)	7476 (7.2)	275 (4.3)	199 (5.0)
Other	8765 (4.6)	414 (3.6)	245 (4.2)	8050 (7.7)	361 (5.6)	250 (6.2)
<i>Missing</i>	5098 (2.7)	255 (2.2)	143 (2.4)	16 (0.02)	0	0
Smoking status - n (%)						
Non-smoker	177 358 (93.3)	10 852 (93.4)	5500 (93.4)	88 263 (84.7)	5465 (85.0)	3403 (84.9)
Smoker	6288 (3.3)	346 (3.0)	167 (2.8)	5205 (5.0)	327 (5.1)	204 (5.1)
<i>Missing</i>	6504 (3.4)	422 (3.6)	224 (3.8)	10 787 (10.4)	639 (10.0)	401 (10.0)
Living with partner - n (%)						
Yes	171 984 (90.4)	10 463 (90.0)	5238 (88.9)	98 403 (94.4)	6042 (94.0)	3749 (93.5)
No	14 047 (7.4)	950 (8.2)	521 (8.8)	4133 (4.0)	278 (4.3)	177 (4.4)
<i>Missing</i>	4119 (2.2)	207 (1.8)	132 (2.2)	1719 (1.7)	111 (1.7)	82 (2.1)

eTable 3. Cont.	Sweden			Norway		
	No hypertension during pregnancy	Hypertension during pregnancy	Preeclampsia	No hypertension during pregnancy	Hypertension during pregnancy	Preeclampsia
Co-existing medical conditions - n (%)						
Gestational diabetes	8345 (4.4)	685 (5.9)	319 (5.4)	6569 (6.3)	612 (9.5)	376 (9.4)
Pre-existing diabetes mellitus	1891 (1.0)	332 (2.9)	206 (3.5)	701 (0.7)	120 (1.9)	90 (2.3)
Chronic kidney disease	655 (0.3)	56 (0.5)	35 (0.6)	555 (0.5)	53 (0.8)	34 (0.9)
Systemic Lupus Erythematosus	267 (0.1)	22 (0.2)	15 (0.3)	81 (0.1)	9 (0.1)	6 (0.2)
History of hypertension during pregnancy - n (%)						
Yes	1248 (0.7)	277 (2.4)	135 (2.3)	2889 (2.8)	827 (12.9)	489 (12.2)
Missing	46 451 (24.4)	2636 (22.7)	1287 (21.8)	0	0	0
Visits to outpatient care 1 year prior						
0 visits	88 962 (46.8)	5170 (44.5)	2610 (44.3)	n/a	n/a	n/a
1-9 visits	52 123 (27.4)	3575 (30.8)	1857 (31.5)	n/a	n/a	n/a
10-29 visits	2566 (1.3)	235 (2.0)	134 (2.3)	n/a	n/a	n/a
30 + visits	48 (0.0)	4 (0.0)	3 (0.1)	n/a	n/a	n/a
Missing	46 451 (24.4)	2636 (22.7)	1287 (21.8)	104 255 (100.0)	6431 (100.0)	4008 (100.0)
Gestational age at birth - n (%)						
Week 22+0 -36+6	7240 (3.8)	1242 (10.7)	1029 (17.5)	4110 (3.9)	811 (12.6)	705 (17.6)
Week 37+0-39+6	86 174 (45.3)	6003 (51.7)	3331 (56.5)	43 165 (41.4)	3222 (50.1)	2103 (52.5)
Week >= 40+0	96 736 (50.9)	4375 (37.7)	1531 (26.0)	56 980 (54.7)	2398 (37.3)	1200 (29.9)

eTable 4. Time between a positive SARS-CoV-2 test and date of diagnosis of hypertension during pregnancy and preeclampsia

Outcome and time (days)	Sweden	Norway
Hypertension during pregnancy		
Mean \pm SD	88.2 \pm 78.5	57.1 \pm 58.0
Median (IQR)	74.0 (24.0-136.0)	45.5 (16.0-82.0)
Range, min-max	-86.0-295.0	-30.0-272.0
Preeclampsia		
Mean \pm SD	86.3 \pm 77.8	60.5 \pm 57.6
Median (IQR)	70.5 (19.0-140.0)	48 (20.0-86.5)
Range, min-max	-77.0-286.0	-20.0-272.0